

PLANTÆ FAVERSHAMIENSES.

A
C A T A L O G U E
OF THE
MORE PERFECT
P L A N T S

Growing spontaneously about

F A V E R S H A M,
IN THE
C O U N T Y O F K E N T.

WITH AN

A P P E N D I X

Exhibiting a SHORT VIEW of the

F O S S I L B O D I E S

OF THE ADJACENT

I S L A N D O F S H E P E Y.

By EDWARD JACOB, Esq; F.S.A.

Consider the Lillies of the Field how they grow — Matt. vii. 28.

L O N D O N:

Printed for the AUTHOR; by J. MARCH, on Tower-Hill.

Sold by B. WHITE, at HORACE'S-HEAD, FLEET-STREET;
and T. EVANS, PATER-NOSTER-ROW.

M,DCC,LXXVII.

14

PLANTAE TAVRSHAMINENSIS.
W. Muggrave!



T O

Mr. JOHN WHITE,
MERCHANT OF LONDON,

IN A GRATEFUL SENSE OF HIS

LONG AND SINGULAR

F R I E N D S H I P.

THIS LITTLE WORK

IS MOST RESPECTFULLY INSCRIBED

BY HIS EVER OBLIGED,

AND VERY HUMBLE SERVANT,

EDWARD JACOB.

Faversham,
Aug. 10, 1777.

M. JOHN WHITE,

MERCHANT OF LONDON,

IN GRATITUDE OF HIS

LONG AND SINGULAR

*Nescio quid fascingerit nostros ut tota mens
irabatur ad peregrinas plantas, illa vero quas
pedibus conculcamus, omnino relinquuntur.*

FROM LINNÆI Bibl. Botan. p. 24.

IS MOST RESPECTFULLY INSCRIBED

BY HIS EVER OBLIGED,

AND VERY HUMBLE SERVANT,

EDWARD JACOB.

P R E F A C E.

THE ensuing Collection of *P L A N T S* was begun many Years since upon the Basis of the late Rev. Mr. Bateman's Catalogue, with whom, in the early part of Life, the Author made several Botanical Excursions hereabouts; and it hath gradually kept increasing ever since, as the constant Visits, in the way of his Profession, to different Parts, gave him opportunities of searching after them. It was ever intended for his own private Use and instructive Entertainment, till lately, at the recommendation of several Gentlemen well versed in Botany, it hath been revised and digested, and is now offered to the Lovers of that Science, as a means, however small, of promoting the topical Knowledge of English Plants; if it shall prove agreeable to their Taste, he will think himself amply rewarded for his Pains.

The great Promoter of its publication was the late worthy Richard Warner, Esq; of Woodford-Row, in Essex, who not only permitted the
free

P R E F A C E.

free Use of his Plantæ Woodfordienses, (without the Aid of which, and Mr. Hudson's excellent Flora Anglica, it would never have been undertaken) but also assisted the Author in the Prosecution of it, and above all, introduced him to the acquaintance of that accurate Botanist, Mr. Alchorne, Assay-Master of the Mint, whose kind Informations and Corrections have been of the greatest Advantage.

The principal Inducement for the Publication was to rectify the Account given in Mr. Blackston's Specimen Botanicum, taken from Mr. Bateman's M. S. Catalogue of near Two Hundred Plants growing about this Town, because many of the rare ones at this time, are not to be found in the Places therein referred to, which have of late been often diligently searched in hope of retrieving them, as they would have added much to this Work, but in vain.

In the Course of about Seventy Years since that Catalogue was made, many Causes and Accidents may be assigned for the loss of some of the rare Plants there mentioned, yet none greater than the industrious hand of improving Husbandry. Mr. Bateman might possibly have been mistaken in others,

P R E F A C E.

others, and possibly too some may be overlooked by the Author, who is not so conceited as to think he hath discovered all the Plants growing round him. Those Gentlemen who are fond of the Study must be sensible, that a single Person, whose necessary attention in the different Branches of his Profession was fully engaged, could not afford sufficient Leisure, however fervent his Inclination might be, to search at proper Seasons every corner of a District Five or Six Miles Diameter, when even the great Mr. Ray could make considerable Additions to the first Catalogue of his Cambridgeshire Plants, and yet leave many unobserved, to be added thereto by Dr. Martyn, and others.

In order, however, to supply his Deficiencies, the Author hath inserted some few rare Plants discovered at a greater distance in the Eastern parts of the County, either by himself, or communicated to him by his obliging Friends, Mr. Alchorne, the Rev. Mr. Jones, of Pluckley, and Mr. Boys of Sandwich, for which Favours he begs they will be pleased to accept his most grateful Acknowledgements.

P R E F A C E.

As he hath composed his Catalogue upon Mr. Warner's alphabetical Plan, he hopes the Preference given to Mr. Ray's Names will be pardoned by the more scientific Botanists, especially as he hath printed an Index of those of Linnæus.

The Appendix, giving a short Account of the Shepey Fossils, it is presumed will not be deemed an improper Addition, by the Lovers of Natural History, as hitherto no particular one hath been offered to their Notice, although so well deserving of it.

The Plate of the Author is the unexpected Donation of an obliging Friend.

THE ABBREVIATIONS explained.

R. S. J. Rail Synopsis Stirpium Britannicarum.
Ed. III. 1724.

H. F. Gul. Hudsoni Flora Anglica. 1762.

A. Annual. B. Biennial. P. Perennial. S. Shrub,
or Tree.

The Month (as March, &c.) denotes the Time of
the Plants flowering.

- * This Mark placed before a Plant, although found wild, shews it most probably to have originated from Seed dispersed by Birds, or otherwise from Gardens.

PLANTÆ FAVERSHAMIENSES.

A

ABSINTHIUM marinum album. R. S. 188.

ARTEMISIA *maritima*. H.F. 311.

White Sea-Wormwood.

In the Salt Marshes—very common. August. P.

ABSINTHIUM vulgare. R. S. 188.

ARTEMISIA *Absinthium*. H.F. 311.

Common Wormwood.

In Farm-yards and on dry banks—common. August. P.

* ACER majus. R. S. 407.

ACER *Pseudo-Platanus*. H.F. 378.

The greater Maple or Sycamore Tree.

In a Hedge near Brenly and Hanslets Forstal — very uncommon. May. S.

ACER minus. R. S. 470.

ACER *campestre*. H. F. 379.

The Common Maple.

In Hedges—very common. May S.

B

ACINOS.

ACINOS. R. S. 238.

THYMUS *Acinos*. H. F. 230.

Wild Basil.

On Badgen Downs — plentifully. July. A.

ADIANTUM *nigrum Officinarum*. R.S. 126.

ASPLENIUM *Adiantum-nigrum*. H.F. 386.

Common black Maiden-hair.

On the hollow sandy Banks in Byfing Wood and
Hernhill—not common. P.

AGRIFOLIUM. R. S. 466.

ILEX *Aquifolium*. H. F. 505.

The Holly Tree.

In Woods and Hedges—common. May. S.

AGRIMONIA. R. S. 202.

AGRIMONIA *Eupatoria*. H. F. 180.

Agrimony.

On the Sides of Fields and under Hedges — very
common. June. S.

ALCEA *vulgaris*. R. S. 252.

MALVA *Alcea*. H. F. 268.

Vervain Mallow.

In Pastures at Graveney and Hernhill — not com-
mon. August. P.

ALLIUM

ALLIUM sylvestre. R. S. 369.

ALLIUM vineale. H. F. 121.

Crow-Garlick.

On the Wall that leads to Thorn Creek — plentifully.
June. P.

ALLIUM sylvestre latifolium. R. S. 370.

ALLIUM ursinum. H. F. 122.

Ramsons.

Amongst the Alders near Hernhill-Church — plentifully. May. P.

ALNUS. R. S. 442.

BETULA Alnus. H. F. 506.

The Common Alder.

In moist Places—very common. July. S.

ALSINE hirsuta Myosotis. R. S. 349.

CERASTIUM vulgatum. H. F. 175.

Narrow-leaved Mouse-ear Chickweed.

In Fields and Meadows—common. May. A.

ALSINE hirsuta Myosotis latifolia præcocior.

R. S. 348.

CERASTIUM viscosum. H. F. 175.

The broader leaved Mouse-ear Chickweed.

On the Walls of the Abbey—common. April. A.

ALSINE major repens perennis. R. S. 347.

CERASTIUM aquaticum. H. F. 177.

Great Marsh Chickweed.

In Davington Brooks, and at Goodnestone — not common. July. P.

ALSINE marina foliis portulacæ. R. S. 351.

ARENARIA peploides. H. F. 167.

Sea Chickweed.

On Shellness in Shepey — not common. July. P.

ALSINE minor multicaulis. R. S. 349.

ARENARIA serpyllifolia. H. F. 167.

The least Chickweed.

On Roofs of old Houses and on old Walls — common.
May. A.

ALSINE spergula dicta major. R. S. 351.

SPERGULA arvensis. H. F. 177.

Corn Spurrey.

On sandy Grounds in Hern-Hill — common. July. A.

ALSINE spergulæ facie media. R. S. 351.

ARENARIA marina. H. F. 169.

Sea Spurrey.

On the Paths to Holly Shore — common. June. A.

ALSINE

ALSINE foliis caryophylleis. R. S. 344.

SAGINA erecta. H. F. 64.

The least Stichwort.

In Nagden Marshes—not common. May. A.

ALSINELLA muscosa flore repens. R. S. 345.

SAGINA procumbens. H. F. 63.

Pin Pearl-wort, Chickweed breakstone.

On old Walls about the Town—common. June. A.

ALTHÆA vulgaris. R. S. 252.

ALTHÆA officinalis. H. F. 267.

Marsh-Mallow.

In the Marshes at Clapgate and Goodnestone—common. August. P.

ANACAMPSEROS, vulgo Faba crassa. R. S. 269.

SEDUM Telephium. H. F. 170.

Orpine, or Live-long.

In Byfing and Cockset Woods—not common.

August. P.

ANAGALLIS flore phoeniceo. R. S. 282.

ANAGALLIS arvensis. H. F. 73.

Male Pimpernel.

In Fields and Gardens—common. May—August. A.

ANAGALLIS lutea. R. S. 282.

LYSIMACHIA nemorum. H. F. 73.

Yellow Pimpernel.

In Jud's Wood, Ospringe—uncommon. June. P.

ANBLATUM Cordi five Aphyllon. R. S. *288.

LATHRÆA Squamaria. H. F. 233.

Toothwort.

Under Hedges near Mr. Jennings's House in Pluckley

—very uncommon. April. P.

ANEMONE nemorum. R. S. 259.

ANEMONE nemorosa. H. F. 208.

Wood Anemony.

In Woods and Hedges—very common. April. P.

ANGELICA sylvestris. R. S. 208.

ANGELICA sylvestris. H. F. 103.

Wild Angelica.

Near Stone-Bridge, and in Byling Wood—not

common. July. P.

• ANGELICA sylvestris minor seu erratica. R. S. 208.

ÆGOPODIUM Podagraria. H. F. 111.

Herb Gerard, Gout-weed, or Ash-weed.

In Gardens and under Hedges, near Boughton Church

—not common. May. P.

ANONIS maritima procumbens. R. S. 332.

ONONIS repens. H. F. 273.

Creeping Rest-harrow.

On the Sand Downs near Deal—uncommon. July. P.

ANONIS

ANONIS spinosa flore purpureo. R. S. 332.

ONONIS spinosa. H. F. 273.

Rest-harrow, Cammock, Petty Whin.

In the Marshes at Clapgate—common. July. P.

ANONIS non spinosa. R. S. 332.

ONONIS arvensis. H. F. 273.

Corn Rest-harrow, or Cammock.

In Corn Fields at Preston and Ospringe—common.
July. P.

ANTIRRHINUM majus. R. S. omittitur.

ANTIRRIHINUM majus. H. F. 239.

The greater Snap-Dragon.

On old Walls about the Town—common. June to
September. P.

APARINE. R. S. 225.

GALIUM Aparine. H. F. 57.

Cleavers, or Goose-grass.

Under Hedges—very common. August. A.

APARINE palustris minor Parisiensis. R. S. 225.

GALIUM uliginosum. H. F. 56.

Marsh Goose-grass.

At the Bottom of Jud's Wood in Ospringe — un-
common. August. P.

APIUM

APIUM palustre et officinarum. R. S. 214.

APIUM graveolens. H. F. 111.

Smallage.

About the Sluice Bridge and the Keys—very common.

August. B.

AQUILEGIA flore simplici. R. S. 273.

AQUILEGIA vulgaris. H. F. 207.

Common Columbines.

In Badgen Wood, and beyond Whitehill, Ospringe

—not common. June. P.

ARTEMISIA vulgaris. R. S. 190.

ARTEMISIA vulgaris. H. F. 312.

Mugwort.

Under Hedges and on the Sides of Fields—very common.

August. P.

ARUM. R. S. 266.

ARUM maculatum. H. F. 342.

Wake-Robin, Cuckow-Pint, Lords and Ladies.

In shady Places and under Hedges—very common.

May. P.

ASPERULA. R. S. 224.

ASPERULA odorata. H. F. 55.

Woodroof.

In Ospringe Woods—plentifully. May. P.

ASPLE-

ASPLENIUM. R. S. 118.

ASPLENIUM *Ceterach*. H. F. 385.

Spleen-wort, Milt-waft.

On Lenham Church Wall — lately lost on the Walls
of our Abbey. P.

ASTER *maritimus cæruleus*. R. S. 175.

ASTER *Tripolium*. H. F. 319.

Sea Star-wort.

In the Salt Marshes — plentifully. August. P.

ASTER *arvensis cæruleus acris*. R. S. 175.

ERIGERON *acre*. H. F. 314.

Blue flowered Flea-bane.

On old Walls and on dry Banks — common. August. P.

ASTER *maritimus flavus*, *Crithmum chrysanthemum dictus*. R. S. 174.

INULA *chrythmoides*. H. F. 320.

Golden Sampire.

In the Marshes of Shepey — uncommon. August. P.

ASTRAGALUS *luteus*. R. S. 326.

ASTRAGALUS *glycyphyllos*. H. F. 281.

Wild Liquorice, or Liquorice Vetch.

Under Hedges in Peasedown, near Whitehill, Ospringe
— uncommon. July. P.

ATRIPLEX *angustifolia maritima dentata.*

R. S. 152.

ATRIPLEX *ferrata.* H. F. 377.

Narrow leaved Orache.

Near Sheerness in Shepey—uncommon. August. A.

ATRIPLEX *angustissimo et longissimo folio.*

R. S. 153.

ATRIPLEX *litoralis.* H. F. 378.

Grass-leaved Orache.

Near Ramsgate Pier in Thanet.—uncommon.

August. A.

ATRIPLEX *maritima.* R. S. 152.

ATRIPLEX *laciniata.* H. F. 376.

Jagged Sea Orache.

Near Sheerness in Shepey—uncommon. August. A.

ATRIPLEX *sylvestris angustifolia.* R. S. 151.

ATRIPLEX *patula.* H. F. 377.

Narrow leaved wild Orache.

In Gardens and on Dunghills—common. August. A.

ATRIPLEX *sylvestris folio hastato seu deltoide.*

R. S. 151.

ATRIPLEX *hastata.* H. F. 377.

Wild Orache with a spear pointed Leaf.

In Gardens and waste Places—common. September. A.

ATRIPLEX

ATRIPLEX *maritima scopariæ folio.* R. S. 153.

ATRIPLEX *litoralis.* H. F. 378. β

Grass-leaved Orache.

Near Margate Pier in Thanet — uncommon.

August. A.

ATRIPLEX *maritima fruticosa.* R. S. 153.

ATRIPLEX *portulacoides.* H. F. 376.

Sea Purslane.

Near Harty Ferry — plentifully. August. P.

ATRIPLEX *maritima perennis folio deltoide seu triangulari.* R. S. 152.

ATRIPLEX *hastata.* H. F. 377. β

Wild Sea Orache.

In the Brent Marshes — common. August. A.

B.

BACCHARIS, *monspeliensium.* R. S. 179.

CONYZA *squarrosa.* H. F. 314.

Plowman's Spikenard.

On Beacon Hill and dry Banks — not uncommon.

August. B.

BALLOTE. R. S. 244.

BALLOTA *nigra.* H. F. 227.

Stinking Horehound.

Under Hedges — very common. August. A.

BELLA-

BELLADONNA. R. S. 265.

ATROPA *Belladonna.* H. F. 79.

Deadly Night-shade, or Dwaile.

Amongst the Buses under Preston-House Wall —

very uncommon. June and July. P.

BELLIS *syvestris minor.* R. S. 184.

BELLIS *perennis.* H. F. 320.

Common Daisy.

In Meadows — very common. From March to October. P.

BERBERIS *dumetorum.* R. S. 465.

BERBERIS *vulgaris.* H. F.

The Berberry-bush, or Pipperidge-bush.

In a Hedge beyond Queen Court, Ospringe — uncommon. May. S.

BETONICA. R. S. 238.

BETONICA *Officinalis.* H. F. 220.

Wood Betony.

In Woods and under Hedges — common. July. P.

BETA *syvestris maritima.* R. S. 38.

BETA *vulgaris.* H. F. 93.

Sea Beet.

On the Sea Walls at Ham and Graveney — uncommon.

August. B.

BETULA.

BETULA. R. S. 443.

BETULA *alba*. H. F. 354.

The Birch Tree.

In Woods—common. April. S.

BIFOLIUM majus, seu Ophrys major quibusdam. R. S. 385.

OPHRYS *ovata*. H. F. 338.

Common Twayblade.

In Cockset and Jud's Woods—common. June. P.

BLITUM Atriplex sylvestris dictum. R. S. 154.

CHENOPODIUM album. H. F. 91.

Common wild Orache.

In uncultivated Places, and on Rubbish—common.

August. A.

BLITUM foetidum Vulvaria dictum. R. S. 156.

CHENOPODIUM Vulvaria. H. F. 92.

Stinking Orache.

Under Walls about the Town—uncommon. August. A.

BLITUM Ficus folio. R. S. 155.

CHENOPODIUM serotinum. H. F. 91.

Late flowering or Fig-leaved Blite.

In Gardens and on Rubbish—common. A.

BLITUM Kali minus album dictum. R. S. 156.

CHENOPODIUM *maritimum*. H. F. 92.

Sea Blite, or white Glass-wort.

In the Salt Marshes—common. August. A.

BLITUM perenne Bonus Henricus dictum.

R. S. 156.

CHENOPODIUM *Bonus Henricus*. H. F. 89.

Common English Mercury, or All-good.

By Way-sides in Ospringe—not uncommon. July. P.

BLITUM Pes anserinus dictum. R. S. 154.

CHENOPODIUM *murale*. H. F. 90.

Common Goose-foot, or Sow-bane.

In Gardens and on Rubbish—not uncommon.
August. A.

BLITUM Pes anserinus dictum acutiore folio.

R. S. 154.

CHENOPODIUM *rubrum*. H. F. 90.

Sharp-leaved Goose-foot, or Sow-bane.

On Dungbills and Rubbish—common. August. A.

BLITUM *rubrum minus*. R. S. 157.

AMARANTUS *Blitum*. H. F. 356.

The least Blite.

On Rubbish and in Gardens—common. August. A.

BORRAGO

BORRAGO hortenſis. R. S. 228.

BORRAGO officinalis. H. F. 68.

Borrage.

On old Walls, and by Road-fides near the Town—
not common. June. P.

BORRAGO ſempervirens. R. S. 227.

ANCHUSA ſempervirens. H. F. 66.

Evergreen Alkanet.

On Rubbiſh near Preſton Church Yard — very un-
common. June. P.

BRASSICA maritima arborea ſive procerior ra-
moſa. R. S. 293.

BRASSICA oleracea. H. F. 253.

Sea Cabbage.

On the Cliffs between Deal and Dover — plentifully.
April. B.

BRYONIA alba. R. S. 261.

BRYONIA alba. H. F. 373.

White Briony.

In Hedges — common. May. P.

BUGLOSSA ſylveſtris minor. R. S. 227.

LYCOPSIS arvenſis. H. F. 69.

Small wild Bugloſs.

In Corn-fields near Preſton Church — uncommon.

May. A.

BUGLOS.

BUGLOSSUM *arvense* annuum. Lithospermifolio. R. S. 227.

LITHOSPERMUM *arvense*. H. F. 66.
Bastard Alkanet.

In Corn-fields in Preston and Hernhill — common
July. A.

BUGULA. R. S. 245.

AJUGA *reptans*. H. F. 219.

Bugle.

In Byfing Wood — common. May. P.

BULBOCASTANUM. R. S. 209.

BUNIUM *Bulbocastanum*. H. F. 105.

Earth-nut, or Hog-nut.

In Byfing Wood and Hernhill — common. June. P.

BUPLEURUM *minimum*. R. S. 221.

BUPLEURUM *tenuissimum*. H. F. 97.

The least Hare's-ear.

Near Sheerness in Shepey — uncommon. August. A.

BUPLEURUM *perfoliatum rotundifolium* annum. R. S. 221.

BUPLEURUM *rotundifolium*. H. F. 97.

Thorow-wax.

In Corn-fields near Brogdale in Ospringe — uncommon.
July. A.

BUTO-

BURSA Pastoris. R. S. 306.

THLASPI *Bursa Pastoris*. H. F. 247.

Shepherd's Purse.

In Hop Grounds, and by Road-sides — very common.

May to September. A.

BUTOMUS. R. S. 273.

BUTOMUS *umbellatus*. H. F. 152.

The flowering Rush, or Water Gladiole.

In Dykes near Pluckley — uncommon. June. P.

* BUXUS. R. S. 445.

BUXUS *sempervirens*. H. F. 355.

The Box Tree.

A few large Trees near Houses in Sheldwich — very uncommon. S.

C.

CAKILE *quibusdam*, aliis *Eruca marina* et *Raphanus marinus*. R. S. 307.

BUNIAS *Cakile*.

Sea Rocket.

Near Sheerness in Shepey, and Cliff's End in Thanet — uncommon. June. A.

CALAMINTHA *humilior folio rotundiore*.

R. S. 243.

GLECHOMA *hederacea*. H. F. 244.

Ground Ivy. Alehoof.

In shady Places, and under Hedges — very common.

May. P.

CALA-

CALAMINTHA vulgaris. R. S. 243.

MELISSA Calamintba. H. F. 230.

Common Calamint.

By the Sides of Roads, and under Hedges — common.

August. P.

CAMPANULA rotundifolia. R. S. 277.

CAMPANULA rotundifolia. H. F. 80.

The lesser round leaved Bell Flower.

By Way-sides in Ospringe and Boughton — not common.

August. P.

CAMPANULA vulgatio foliis urticæ. R. S. 276.

CAMPANULA Trachelium. H. F. 81.

Canterbury Bells, or Great Throat-wort.

In Woods and Hedges — very common. — With a white

Flower in Hernhill — very uncommon. July. P.

CAPRIFOLIUM germanicum. R. S. 458.

LONICERA Periclymenum. H. F. 80.

Common Honey-suckle, or Woodbine.

In Woods and Hedges — very common. April to

August. S.

CARDAMINE. R. S. 299.

CARDAMINE pratensis. H. F. 256.

Common Ladies Smock, or Cuckow-flower.

In moist Meadows — very common. April. P.

CARDI-

CARDAMINE parviflora. H. F. 257.

Small flowered Ladies Smock.

In Meadows near Pluckley—*uncommon*. April. A.

CARDIACA. R. S. 239.

LEONURUS Cardiac. H. F. 228.

Mother-wort.

Near Cockset, in Ospringe — *very uncommon*.

July. B.

CARDUUS acaulos minore purpureo flore.

R. S. 195.

CARDUUS acaulas. H. F. 308.

Dwarf Carline Thistle.

Upon Beacon Hill—*not uncommon*. July. P.

CARDUUS caule crispo. R. S. 194.

CARDUUS crispus. H. F. 306.

Thistle upon Thistle.

By the Road-sides, and on dry Banks in Ospringe —
common. June. A.

CARDUUS lanceolatus. R. S. 195.

CARDUUS lanceolatus. H. F. 305.

Spear-Thistle.

In uncultivated Ground, and under Hedges — *common*.

July. B.

CAR-

CARDUUS Mariæ. R. S. 195.

CARDUUS *marianus*. H. F. 307.

Milk Thistle, or Lady's Thistle.

*By the Way-side at Davington, and by the Sea Walls
beyond Nagden—not uncommon. July. A.*

CARDUUS Mariæ *hirsutus non maculatus*.

R. S. 194.

CARDUUS *marianus*. H. F. 307. β.

Ladies Thistle without Spots.

In Graveney Marshes — uncommon. July. A.

CARDUUS *nutans*. R. S. 193.

CARDUUS *nutans*. H. F. 306.

Musk Thistle.

On Badgen Downs — plentifully. July. B.

CARDUUS *palustris*. R. S. 194.

CARDUUS *palustris*. H. F. 306.

Marsh Thistle.

In the Marshes — common. July. B.

CARDUUS *spinosissimus capitulis minoribus*.

R. S. 194.

CARDUUS *acanthoides*. H. F. 306.

Wetted Thistle with small Flowers.

Upon Graveney Marsh Sea Wall—common. July. A.

CAR-

CARDUUS stellatus. R. S. 176.

CENTAUREA Calcitrapa. H. F. 326.

Star Thistle.

In Graveney Marshes — common. July. A.

CARDUUS tomentosus **Acanthium** dictus vulgaris. R. S. 196.

ONOPORDUM **Acanthium**. H. F. 308.

Cotton Thistle.

In uncultivated Places, and by the Road-side — common.

July. B.

CARDUUS vulgatissimus viarum. R. S. 194.

SERRATULA arvensis. H. F. 305.

Common Way-Thistle, or Creeping Thistle.

In Fields and Byways — very common. July. P.

CARLINA sylvestris quibusdam, aliis **Atractilis**.

R. S. 175.

CARLINA vulgaris. H. F. 309.

The common wild Carline Thistle.

In dry Fields and Pastures — common. June. B.

CARYOPHYLLATA. R. S. 253.

GEUM urbanum. H. F. 198.

Herb Bennet, or Common Avens.

In Woods, and under Hedges — common. July. P.

CARY

CARYOPHYLLATA montana purpurea. R. S.
253.

GEUM rivale. H. F. 198.

Water Avens.

In a Wood near Barber's Mill in Hothfield — very
uncommon. July. P.

CARYOPHYLLUS holosteus arvensis glaber flore
majore. R. S. 346.

STELLARIA Holostea. H. F. 166.

The greater Stich-wort.

Under Hedges, and on the Sides of Fields — very
common. April. P.

CARYOPHYLLUS holosteus arvensis glaber flore
minore. R. S. 346.

STELLARIA graminea. H. F. 166.

The lesser Stich-wort.

Under Hedges, chiefly on a dry Soil — common. July. P.

CARYOPHYLLUS holosteus arvensis medius.
R. S. 347.

STELLARIA graminea. H. F. 166. β .

The middle Stich-wort.

Upon Beacon Hill — not uncommon. July. P.

CARYOPHYLLUS latifolius barbatus minor
annuus flore minore. R. S. 337.

DIANTHUS Armeria. H. F. 161.

Deptford Pink.

On Beacon Hill, and dry Pastures in Ospringe — not
uncommon. July. A. CAS-

R. S. CASTANEA. R. S. 440.

FAGUS *Castanea*. H. F. 359.

The Chesnut Tree.

Byfing and other Woods—not uncommon. May. S.

very
EATANANCE leguminosa quorundam. R. S.
325.

flor
LATHYRUS *Nissolia*. H. F. 275.

Crimson Grass-Vetch.

On Beacon Hill — not uncommon. June. A.

very
CAUCALIS minor flosculis rubentibus. R. S. 219.

CAUCALIS *Anthriscus*. H. F. 99.

flor
Hedge Parsley.

In bushy Grounds, and on Ham Wall—not uncommon.
July. B.

P.
CAUCALIS segetum minor, Anthriscus hispido
similis. R. S. 220.

is.
CAUCALIS *arvensis*. H. F. 98.

Small Corn Parsley.

In Fields, and amongst Corn—common. August. A.

CENTAURIUM luteum perfoliatum. R. S. 287.

BLACKSTONIA *perfoliata*. H. F. 146.

Yellow Centory.

In dry chalky Fields — common. July. A.

CEN-

not
S-

CENTAURIUM minus. R. S. 286.

GENTIANA Centaurium. H. F. 88.

Lesser Centory.

In dry Pastures—very common. July to October. A.

*With a white Flower, near Minster in Shepey —
but uncommon.*

CERASUS sylvestris fructu rubro. R. S. 463.

PRUNUS Avium. H. F. 187.

Common wild Cherry.

In Woods — common. April. S.

CERASTIUM hirsutum minus parvo flore.

R. S. 348.

CERASTIUM semi-decandrium. H. F. 175.

The least Mouse-ear Chickweed.

On Walls and dry Soils — common. April. A.

CEREFOLIUM sylvestre. R. S. 207.

CHÆROPHYLLUM temulum. H. F. 108.

Wild Chervil.

Under Hedges — very common. July. A.

CHAMÆMELUM fætidum. R. S. 185.

ANTHEMIS Cotula. H. F. 223.

Stinking Mayweed.

In Corn Fields — too common. June. A.

CHA-

CHAMÆMELUM inodorum. R. S. 185.

MATRICARIA inodora. H. F. 322.

Field Feverfew.

On the sides of paths in Corn-fields — common.
August. A.

CHAMÆMELUM odoratissimum repens flore
simplici. R. S. 185.

ANTHEMIS nobilis. H. F. 323.

Sweet-scented Camomile.

On Charing Heath — uncommon. August. P.

CHAMÆMELUM vulgare. R. S. 184.

MATRICARIA Chamomilla. H. F. 322.

Corn Feverfew.

In cultivated Fields amongst Corn — common. June. A.

CHELIDONIUM minus. R. S. 246.

FICARIA verna. H. F. 214.

Pilewort, or lesser Celandine.

In Meadows — very common. April. P.

CHENOPODIUM Betæ folio. R. S. 157.

CHENOPODIUM polyspermum. H. F. 92.

Upright round leaved Blite, or Allseed.

In the Conduit Row — not uncommon. August. A.

D

CHENO.

CHENOPODIUM erectum foliis triangularibus
dentatis, spicis e foliorum alis plurimis longis
erectis tenuibus. R. S. 155.

CHENOPODIUM urbicum. H. F. 89.

Upright Blite.

In the Conduit Row, and in Ditches—common. Sep-
tember. A.

CHENOPODIUM Stramonii folio. R. S. 154.

CHENOPODIUM hybridum. H. F. 90.

Maple-leaved Blite.

On Dungbills—not common. August. A.

CHRYSANTHEMUM segetum. R. S. 182.

CHRYSANTHEMUM segetum. H. F. 321.

Corn Marigold.

In Corn-fields—common. July. A.

CICHOREUM fylvestre. R. S. 172.

CICHOREUM Intybus. H. F. 303.

Wild Succory.

*In uncultivated Places, and by Sides of Fields—very
common.* August. P.

CICUTA. R. S. 215.

CONIUM maculatum. H. F. 100.

Hemlock.

In shady Places near the Town—common. July. A.

CICU-

CICUTARIA tenuifolia. R. S. 215.

ÆTHUSA *Cynapium*. H. F. 107.

The lesser Hemlock, or Fools Parsley.

In Gatefield, and in Gardens — not uncommon. September. A.

CICUTARIA vulgaris. R. S. 207.

CHÆROPHYLLUM *sylvestre*. H. F. 108.

Wild Cicely, or Cow-weed.

Under Hedges, and in Fields — common. June. A.

CIRCÆA lutetiana. R. S. 289.

CIRCÆA *lutetiana*. H. F. 9.

Enchanters Night-shade.

In Byfing Wood — common. July. P.

CLEMATIS daphnoides major. R. S. 268.

VINCA *major*. H. F. 77.

The greater Periwinkle.

In Hedges, near Houses at Ore and Sindal Farm — uncommon. April. P.

CLEMATIS latifolia feu Atragene quibusdam.
R. S. 258.

CLEMATIS *Vitalba*. H. F. 217.

Great Wild Climber, or Traveller's Joy.

In Hedges — very common. July. P.

CLINOPODIUM *Origano simile.* R. S. 239.

CLINOPODIUM *vulgare.* H. F. 228.

Great wild Basil.

In Hedges — very common. July. P.

COCHLEARIA folio sinuato. R. S. 303.

COCHLEARIA *anglica.* H. F. 248.

Sea Scurvy-grass.

On the Sea Walls near Thorn — common. May. B.

CONVOLVULUS major. R. S. 275.

CONVOLVULUS *sepium.* H. F. 74.

Great Bindweed.

Under Hedges — very common. With a reddish purple Flower, in the Brent Hedge; but very uncommon.
August. P.

CONVOLVULUS *maritimus Soldanella dictus.*
R. S. 276.

CONVOLVULUS *Soldanella.* H. F. 75.

Sea Bindweed.

On the Sand Downs near Sandwich — not common.
July. P.

CONVOLVULUS *minor vulgaris.* R. S. 275.

CONVOLVULUS *arvensis.* H. F. 74.

Small Bindweed.

In Fields, and by Road-sides — very common. July. P.

CONIZA

CONYZA media. R. S. 174.

INULA dysenterica. H. F. 320.

Middle Fleabane.

About the Abbey Pond — common. August. P.

CONYZA minor. R. S. 174.

INULA Pulicaria. H. F. 320.

Small Fleabane.

About the Conduit-row, and at Downe's-forstal in
Hernhill — not uncommon. September. A.

CORNUS fæmina. R. S. 460.

CORNUS sanguinea. H. F. 58.

The female Cornel, Dog-berry Tree, or
Prick-wood.

In Hedges — common. June. S.

CORYLUS sylvestris. R. S. 439.

CORYLUS Avellana. H. F. 360.

The Hazle-nut Tree.

In Woods and Hedges — very common. March. S.

COTYLEDON vera radice tuberosa. R. S. 271.

COTYLEDON Umbilicus Veneris. H. F. 169.

Navel-wort, Kidney-wort, or Wall Penny-wort.

On Tenterden Church, and in a Stone Pit at
Boughton Monchelsea — very uncommon. June. P.

CRACCA. R. S. 322.

VICIA *Cracca*. H. F. 277.

Tufted Vetch.

In the Hedges of the Beacon Farm — uncommon.

July. A.

CRACCA minor. R. S. 322.

ERVUM *hirsutum*. H. F. 280.

Small wild Tare, Tine-Tare, or hairy Tare.

In Rickefy-lane Hedge, Selling — not uncommon.

July. A.

CRACCA minor filiquis singularibus flosculis
cærulescentibus. R. S. 322.

ERVUM *tetraspermum*. H. F.

Tine-Tare with smooth Pods, or smooth Tare.

In Fields and under Hedges — common. June. A.

CRITHMUM *maritimum*. R. S. 217.

CRITHMUM *maritimum*. H. F. 101.

Sampire.

*On the Cliffs between St. Margaret's and Dover —
plentifully. August. P.*

CRUCIATA. R. S. 223.

VALANTIA *Cruciata*. H. F. 375.

Cross-wort, or Mug-weed.

In Hedges in Preston and Ospringe — not uncommon.

June. F.

CUSCUTA

CUSCUTA major. R. S. 281.

CUSCUTA *europæa*. H. F. 89.

Dodder, Hell-weed, Devil's-guts.

Upon leguminous Plants — common. July. A.

CYANUS. R. S. 198.

CENTAUREA *Cyanus*. H. F. 325.

Blue Bottles, or Corn Bottles.

In Corn-fields — very common. July. A.

CYNOGLOSSUM. R. S. 226.

CYNOGLOSSUM *officinale*. H. F. 67. β.

Great Hound's Tongue.

By the Road Sides near Boughton Chalk Pits — ~~not~~
common. June. P.

CYNOGLOSSUM *folio-virentia*. R. S. 226.

CYNOGLOSSUM *officinale*. H. F. 67. β.

The lesser green leaved Hound's Tongue.

Near Sandwich — not common. June. P.

D.

DAMASONIUM *stellatum* Dalechampii.

R. S. 272.

ALISMA *Damasonium*. H. F. 138.

Star-headed Water Plantain.

In a Field-pond of Mr. Jacob's Farm, at East Church
in Shepey — very uncommon. June to August. P.

DAU.

DAUCUS vulgaris. R. S. 218.

DAUCUS *Carota*. H. F. 99.

Wild Carrot, or Bird's Nest.

In Pastures, and by Way-sides—very common. July. B.

• DELPHINIUM *segetum flore caruleo*.

R. S. 273.

DELPHINIUM *Consolida*. H. F. 207.

Wild Larkspur.

In King's Field, amongst Corn — very uncommon.

June. A.

DENS-LEONIS. R. S. 170.

LEONTODON *Taraxacum*. H. F. 297.

Dandelion.

In Meadows and Pastures—very common. May. P.

DENS-LEONIS *hirsutus*, *Hieracium dictus*.

R. S. 127.

LEONTODON *hispidum*. H. F. 297.

Rough Dandelion, or Dandelion Hawk-weed.

In Meadows and Pastures—common. June. P.

DIGITALIS *purpurea*. R. S. * 283.

DIGITALIS *purpurea*. H. F. 240.

Purple Fox Glove.

In Woods and Hedges—common. July. B.

DIPSACUS

DIPSACUS minor seu Virga pastoris. R. S. 192.

DIPSACUS pilosus. H. F. 49.

Small wild Teasel, or Shepherd's Rod.

In the Road Hedge leading from Plomford to Badgen

Downs — uncommon. July or August. B.

DIPSACUS sylvestris seu Labrum Veneris.

R. S. 192.

DIPSACUS sylvestris. H. F. 49.

Wild Teasel.

In uncultivated Places, and under Hedges — very common. July. B.

E.

ECHIUM alterum sive Lycopsis anglica. R. S. 228.

ECHIUM anglicum. H. F. 70.

English Viper's Bugloss.

By Way sides near Whitehill in Ospringe — not uncommon. August. P.

ECHIUM vulgare. R. S. 218.

ECHIUM vulgare. H. F. 69.

Viper's Bugloss.

On barren Ground, and upon old Walls — very common.

July. P.

EQUISETUM arvense longioribus setis, R. S.

130.

EQUISETUM arvense. H. F. 380.

Corn Horse-tail.

In moist Fields amongst Corn — common. March. P.

EQUISETUM majus. R. S. 130.

EQUISETUM fluviatile. H. F. 381.

River Horse-tail.

In the River about the Powder Mills — common.

May. P.

EQUISETUM palustre. R. S. 131.

EQUISETUM palustre. H. F. 380.

The lesser Marsh Horse-tail.

In the Corn Fields near Nagden — not uncommon.

June. P.

EQUISETUM sylvaticum. R. S. 130.

EQUISETUM sylvaticum. H. F. 380.

Wood Horse-tail.

In the swampy Ground near Ore Mill — not common.

May. P.

ERICA Brabantica folio Coridis hirsuto quaterno.

R. S. 471.

ERICA tetralix. H. F. 144.

Cross-leaved Heath.

At the Bottom of Jud's Wood — not common. Au-

gust. S.

ERICA tenuifolia. R. S. 471.

ERICA cinerea. H. F. 144.

Fine leaved Heath.

Upon Charing Heath — not common. August. S.

ERICA

ERICA vulgaris. R. S. 470.

ERICA vulgaris. H. F. 144.

Common Heath, or Ling.

In Woods—very common. June to September. P.

ERUCA aquatica. R. S. 297.

SISYMBRIUM sylvestre. H. F. 258.

Water Rocket.

In Ham Ponds near Sandwich—uncommon. June. P.

ERUCA hirsuta filiqua cauli appressa. Erysimum
vulgare. R. S. 298.

ERYSIMUM officinale. H. F. 250.

Hedge Mustard.

Under Hedges, and by Road Sides — very common.
May. A.

ERUCA lutea seu Barbarea. R. S. 297.

ERYSIMUM Barbarea. H. F. 251.

Winter Cress or Rocket.

Near the Powder Mills—not uncommon. May. P.

ERYNGIUM marinum. R. S. 222.

ERYNGIUM maritimum. H. F. 95.

Eryngo, or Sea Holly.

On the Shore between Graveney and Sea Salter —
uncommon. July. P.

ERYSIMUM

ERYSIMUM Sophia dictum. R. S. 298.

SYSIMBRIUM Sophia. H. F. 259.

Flix-weed.

By the Road Sides near Sittingbourn — uncommon.

June. A.

EUONYMUS vulgaris. R. S. 468.

EUONYMUS europæus. H. F. 84.

The Spindle Tree, or Prick-wood.

In Woods and Hedges — common. May. S.

EUPATORIUM Cannabinum. R. S. 179.

EUPATORIUM cannabinum. H. F. 310.

Common Hemp Agrimony, or Dutch Agrimony.

About the Abbey Pond — common. August. P.

EUPHRASIA. R. S. * 284.

EUPHRASIA officinalis. H. F. 234.

Eye-bright.

In most upland Pastures — very common. August. A.

EUPHRASIA pratensis rubra. R. S. * 284.

EUPHRASIA Odontites. H. F. 234.

Red Eye-bright.

In the Ways and Corn Fields about Preston — very
common. August. A.

FAGUS.

F.

FAGUS. R. S. 439.

FAGUS *sylvaticus*. H. F. 360.

The Beech Tree.

In Woods — common. May. S.

* FEGOPYRUM. R. S. 144.

POLYGONUM *Fagopyrum*. H. F. 150.

Buck-wheat, or Brank.

In Corn fields near Beacon Hill — uncommon. July. A.

FEGOPYRUM *scandens sylvestre*. R. S. 144.

POLYGONUM *Convolvulus*. H. F. 149.

Black Bind-weed.

In Corn fields and Hedges — not uncommon. June to
September. A.

FERRUM *equinum germanicum filiquis in sum-*
mitate. R. S. 326.

HIPPOCREPIS *comosa*. H. F. 281.

Tufted Horse-shoe Vetch.

On chalky Ground in Ospringe — common. July. A.

FILIPENDULA. R. S. 259.

SPIRÆA *Filipendula*. H. F. 190.

Common Dropwort.

Upon Beacon Hill — not uncommon. July. P.

E

FILIX

FH IX Fœmina. R. S. 124.

PTERIS *aquilina*. H. F. 384.

Female Fern, or Common Brakes.

In Woods — very common. August. P.

FILIX mas vulgaris. R. S. 120.

POLYPODIUM *Filix mas*. H. F. 389.

Common Male Fern.

On shady Banks under Hedges — common. P.

FILICULA saxatilis ramosa maritima nostras.

R. S. 125.

PTERIS *aquilina*. H. F. 384. β .

Small-branched Stone-Fern.

On Walls about the Town — uncommon. August. P.

FCENICULUM vulgare. R. S. 217.

ANETHUM *Fœniculum*. H. F. 110.

Common Fennel, or Finkle.

By the Road-side as you enter the Town — not uncommon. August. P.

FRAGARIA. R. S. 254.

FRAGARIA *vesca*. H. F. 194.

Strawberry.

In Woods — very common. April. P.

FRAGRA-

FRAGARIA *sterilis*. R. S. 254.

FRAGARIA *sterilis*. H. F. 195.

Barren Strawberry.

On barren Lands — very common. April. P.

FRANGULA *sive Alnus nigra baccifera*. R.
465.

RHAMNUS *Frangula*. H. F. 83.

The black Berry-bearing Alder.

In Badgen Wood — not common. May. S.

FRAXINUS. R. S. 469.

FRAXINUS *excelsior*. H. F. 379.

The common Ash Tree.

In Woods and Hedges — common. April. S.

FUMARIA *vulgaris*. R. S. 204.

FUMARIA *officinalis*. H. F. 270.

Fumitory.

In Fields and Gardens — common. May. A.

FUMARIA *alba latifolia*. R. S. 204.

FUMARIA *claviculata*. H. F. 270.

Climbing Fumitory.

*In moist Hedges in Davington and Ospringe — not
common. August. A.*

G.

GALE frutex odoratus septentrionalium. R. S. 443.

MYRICA Gale. H. F. 368.

Sweet Willow, Goule, Dutch Myrtle.

On Willsborough Lees, near Ashford — plentifully.

May. S.

GALEOPSIS legitima Dioscoridis. R. S. 237.

STACHYS sylvatica. H. F. 227.

Hedge Nettle.

In Woods and Hedges—very common. August. A.

GALLIUM luteum. R. S. 224.

GALLIUM verum. H. F. 55.

Yellow Ladies Bedstraw, or Cheese-renning.

On the Sides of Fields—very common. July. P.

GENISTA angulosa trifolia. R. S. 474.

SPARTIUM scoparium. H. F. 271.

Common Broom.

In Woods and dry Fields — very common. June. P.

GENISTA minor aspalathoides. R. S. 475.

GENISTA anglica. H. F. 272.

Needle Furze, or Petty Whin.

Amongst Furze which grows on wet Land — not common. May. P.

GENISTA

GENISTA spinosa vulgaris. R. S. 475.

ULEX europæus. H. F. 272.

Furze, Whins, or Gorse.

On Beacon Hill—very common. March to June. P.

GENISTELLA tinctoria. R. S. 474.

GENISTA tinctoria. H. F. 272.

Dyers-weed, Green-wood.

In the Marshes near Holly Shore — not common.

July. P.

GENTIANELLA fugax autumnalis. R. S. 275.

GENTIANA Amarella. H. F. 87.

Autumnal Gentian.

In dry Pastures and Ofpringe Chalk Pits — not uncommon. August. A.

GERANIUM Cicutæ folio inodorum. R. S. 357.

GERANIUM cicutarium. H. F. 262.

Field Crane's-bill without Scent, or Hemlock-leaved Crane's-bill.

By Road sides, and on old Walls — not uncommon.

April to June. A.

GERANIUM columbinum. R. S. 359.

GERANIUM molle. H. F. 265.

Common Dove's-foot Crane's bill.

Under Hedges — very common. June. A.

GERANIUM columbinum humile flore cæruleo.

R. S. 359.

GERANIUM pusillum. H. F. 266.

Small flowered Dove's-foot Crane's-bill.

By Waysides leading to Thorn—common. June. A.

GERANIUM columbinum majus foliis dissectis.

R. S. 359.

GERANIUM dissectum. H. F. 266.

Jagged leaved Dove's-foot Crane's-bill.

In dry Fields in Ospringe — not uncommon. May to July. A.

GERANIUM columbinum dissectis foliis pediculis

florum longissimis. R. S. 359.

GERANIUM columbinum. H. F. 266.

Long-stalked Dove's-foot Crane's-bill.

Upon Oldwives Lees — not common. July. A.

GERANIUM robertianum. R. S. 358.

GERANIUM robertianum. H. F. 264.

Herb Robert.

In Hedge, and on Rubbish—very common. June. B.

GERANIUM pusillum supinum maritimum.

R. S. 357.

GERANIUM maritimum. H. F. 263.

Sea Crane's-bill.

On the Sand Downs near Deal — not common.

August. P.

GLAUX

GLAUX *maritima*. R. S. 285.

GLAUX *maritima*, H. F. 86.

Sea Milkwort, or black Saltwort.

In the Brent Marshes — common. July. P.

GNAPHALIUM *longifolium humile ramosum*
capitulis nigris. R. S. 181.

GNAPHALIUM *uliginosum*. H. F. 313.

Black-headed, long leaved, low branched cud-
weed.

In moist Places in Hernhill — not uncommon.
August. A.

GNAPHALIUM *anglicum*. R. S. 180.

GNAPHALIUM *sylvaticum*. H. F. 313.

Upright Cudweed.

In Judd's Wood — common. August. B.

GNAPHALIUM *minimum*. R. S. 181.

FILAGO *montana*. H. F. 328.

The least Cudweed.

In dry sandy Meadows, and among Corn — common.
July. A.

GNAPHALIUM *minus* feu *Herba impia*.
R. S. 180.

FILAGO *germanica*. H. F. 328.

Common Cudweed.

In dry Meadows and by Road sides — common. July. A.

GROS-

* GROSSULARIA. Miller Botan. Officinal. 220.

RIBES *Uva crispa*. Lin. System. Nat. Tom. II.

p. 184.

The Gooseberry Bush.

In several Hedges beyond White Hill, Ospringe.
April. S.

H.

HEDERA communis major et minor. R. S. 459.

HEDERA *Helix*. H. F. 85.

Climbing, or berried Ivy, also barren and creeping
Ivy.

In Woods and Hedges — very common. October. S.

HELIANTHEMUM vulgare. R. S. 341.

CISTUS *Helianthemum*. H. F. 205.

Dwarf Cistus, or little Sunflower.

On dry chalky Banks — very common. June. P.

HELENIUM. R. S. 176.

INULA *Helenium*. H. F. 319.

Elecampane.

*On the moist Ground near the Half-way House to Can-
terbury — very uncommon.* August. P.

HELLEBORUS niger hortensis. R. S. 271.

HELLEBORUS *viridis*. H. F. 215.

Wild black Hellebore.

*On the Cliff beyond Westfield in Pluckley — un-
common.* April. P. HEL.

HELLEBORASTER maximus. R. S. 271.

HELLEBORUS *fætidus*. H. F. 215.

Great bastard black Hellebore, Bear's-foot, or
Setter-wort.

By the Road side, up the Chalk-hill, about a Mile
North-west from Charing—*uncommon*. March. P.

HELLEBORINE *altera atro-rubente flore*.
R. S. 383.

SERAPIAS *latifolia*. H. F. 341. β .

Broad-leaved bastard Hellebore, a Variety.

In Ospringe Woods—*uncommon*. August. P.

HELLEBORINE *latifolia montana*. R. S. 383.

SERAPIAS *latifolia*. H. F. 341.

Broad leaved bastard Hellebore.

In King's Wood — *uncommon*. July. P.

HELLEBORINE *latifolia flore albo clauso*.
R. S. 384.

SERAPIAS *longifolia*. H. F. 341.

White flowered bastard Hellebore.

In Badgen and Jud's Woods—*uncommon*. June. P.

HERBA Paris. R. S. 264.

PARIS *quadrifolia*. H. F. 150.

Herb Paris, True-love, or One-berry.

In Woods near Doddington—*not common*. May. P.

HES.

HESPERIS *allium redolens.* R. S. 293.

ERYSIMUM *Alliaria.* H. F. 251.

Jack by the Hedge, or Sauce alone.

Under Hedges — very common. May. P.

HIERACIUM *Echioides capitulis Cardui benedicti.* R. S. 166.

PICRIS *Echioides.* H. F. 294.

Ox's Tongue, or Lang de Boeuf.

In Hedges and Pastures going to Thorn Creek — common. July. A.

HIERACIUM *fruticosum angustifolium majus.* R. S. 168.

HIERACIUM *umbellatum.* H. F. 300.

Narrow leaved bushy Hawk-weed.

In Ospringe Chalk Pits, and by Way-sides at Dunkirk — uncommon. August. P.

HIERACIUM *fruticosum latifolium hirsutum.* R. S. 187.

HIERACIUM *sabaudum.* H. F. 300.

Bushy Hawk-weed with broad rough Leaves.

In Hedges about Sheldwich — not common. July. P.

HIERACIUM *longius radicum.* R. S. 165.

HYPOCHÆRIS *radicata.* H. F. 302.

Long-rooted Hawk-weed.

On Beacon Hill — common. June. P.

HIER.

HIERACIUM luteum glabrum five minus hirsutum. R. S. 165.

CREPIS testorum. H. F. 301.

Smooth Succory Hawk-weed.

In Fields and on old Walls — common. June to September. A.

HIERACIUM maximum Chondrillæ folio asperum. R. S. 166.

CREPIS biennis. H. F. 301.

* Rough Succory Hawk-weed.

In Hedges near Sittingbourn — uncommon. August. B.

HIERACIUM minus præmorsa radice. R. S. 164.

LEONTODON autumnale. H. F. 297.

Hawk-weed with bitten Roots, or yellow Devil's Bit.

In Corn fields in Hernhill — common. August. P.

HORMINUM sylvestre Lavendulæ flore. R. S. 237.

SALVIA verbenaca. H. F. 9.

Common English wild Clary, or Oculus Christi.

In Davington and Ore Church-Yards — not common.

May. P.

HOTTONIA. R. S. 285.

HOTTONIA palustris. H. F. 72.

Water Violet, or Gilliflower.

About the Decoy Ponds — uncommon. *In Dykes near*

Deal — common. June. P.

HYA

HYACINTHUS anglicus. R. S. 373.

HYACINTHUS *non scriptus*. H. F. 123.

English Hyacinth, or Hare-bells.

In Woods and under Hedges—very common. April. P.

HYDROCOTYLE vulgaris. R. S. 222.

HYDROCOTYLE vulgaris. H. F. 96.

Marsh Penny-wort, or White-rot.

*In the Marshes about Luddenham and Ore — not
uncommon.* May. P.

HYOSCYAMUS vulgaris. R. S. 274.

HYOSCYAMUS *niger*. H. F. 77.

Common Henbane.

By Road sides near the Town—common. June. P.

HYPERICUM. R. S. 342.

HYPERICUM *perforatum*. H. F. 290.

Saint John's-wort.

Under Hedges, and by Road sides — very common.
July. P.

HYPERICUM *Androsæmum dictum*. R. S. 343.

HYPERICUM *hirsutum*. H. F. 291.

Tutsan, or hairy St. John's-wort.

In hollow Lanes in Boughton—uncommon. July. P.

HYPE-

HYPERICUM Ascyron dictum caule quadrangulo. R. S. 344.

HYPERICUM *quadrangulum*. H. F. 292.

Saint Peter's-wort.

In Davington Brooks—not uncommon. July. P.

HYPERICUM pulchrum Tragi. R. S. 342.

HYPERICUM *pulchrum*. H. F. 290.

Upright Saint John's-wort.

In Byfing Wood — common. July. P.

HYPERICUM maximum Androsæmum vulgare dictum. R. S. 343.

HYPERICUM *Androsæmum*. H. F. 291.

Tutfan, or Park-leaves.

In a Hedge near Provender Wood — uncommon.

July. P.

HYPERICUM minus supinum. R. S. 343.

HYPERICUM *humifusum*. H. F. 299.

The least trailing Saint John's-wort.

On sandy Lands in Ore—common. July. P.

HYPERICUM *elegantissimum non ramosum folio lato*. R. S. 343.

HYPERICUM *montenum*. H. F. 291.

Imperforate Saint John's-wort.

In Byfing Wood — uncommon. July. P.

J.

JACEA major. R. S. 198:

CENTAUREA *Scabiosa*. H. F. 325.

Great Knap-weed, or Matfellow.

On the sides of Paths — common. July. P.

JACEA nigra. R. S. 198.

CENTAUREA *Jacea*. H. F. 326.

Common Knap-weed, or Matfellow.

On the sides of Fields — very common. August. P.

JACOBÆA latifolia palustris five aquatica.

R. S. 178.

SENECIO *aquaticus*. H. F. 317.

Broad-leaved Marsh or Water Ragwort.

In the Brents — common. August. P.

JACOBÆA senecionis folio incano perennis.

R. S. 177.

SENECIO *erucifolius*. H. F. 317.

Hoary Perennial Ragwort with Groundsel Leaves.

In Byfing Wood — not common. July. P.

JACOBÆA vulgaris. R. S. 177.

SENECIO *Jacobæa*. H. F. 316.

Common Ragwort.

In Meadows and Pastures — very common. July. P.

IRIS

IRIS palustris lutea. R. S. 374.

IRIS Pseudacorus. H. F. 13.

Yellow Water Flower de luce.

In moist boggy Places — very common. June. P.

IRIS sylvestris quam Xyrim vocant. R. S. 375.

IRIS fœtidissima. H. F. 13.

Stinking Gladdon.

By Sandgate Castle near Folkstone — very uncommon.

July. P.

JUNIPERUS vulgaris baccis parvis purpureis.

R. S. 444.

JUNIPERUS communis. H. F. 372.

The common Juniper.

On dry chalky Commons — very common. May. S.

K.

KALI geniculatum perenne fruticosius procumbens. R. S. 136.

SALICORNIA europæa. H. F. 1. β.

Perennial Marsh Sampire, or jointed Glass-wort.

In the Isle of Shepey — uncommon. August. P.

KALI spinosum cochleatum. R. S. 159.

SALSOLA Kali. H. F. 33.

Prickly Glass-wort.

On the Shore between Graveney and Seasalter — not common. July. A.

KNAWEL. R. S. 159.

SCLERANTHUS *annuus*. H. F. 160.

German Knot-grass, or annual Knawel.

On barren dry sandy Soils—very common. August. A.

L.

LACTUCA *agnina* seu *Valerianella foliis ferratis*.

R. S. 201.

VALERIANA *Locusta*. H. F. 12.

Small Corn Sallet, or Valerian with jagged leaves.

In Fields among Corn, and on old Walls——common.

May. A.

LACTUCA *sylvestris murorum*. R. S. 162.

PRENANTHES *muralis*. H. F. 296.

Ivy-leaved wild Lettuce.

In Hedges near Wilderton—uncommon. July. P.

LACTUCA *sylvestris minima*. R. S. 162.

LACTUCA *saligna*. H. F. 296.

The least wild Lettuce.

On Banks of Ditches near Sheerness——uncommon.

August. A.

LAMIUM *album*. R. S. 240.

LAMIUM *album*. H. F. 225.

White Archangel or Dead Nettle.

Under Hedges——very common. May. P.

LAMIUM

LAMIUM rubrum. R. S. 240.

LAMIUM rubrum. H. F. 225.

Red Archangel or Dead Nettle.

On Rubbish, and sides of Fields — very common.

May. A.

LAMIUM cannabinum floribus albis verticillis
purpurascentibus. R. S. 241.

GALEOPSIS *Tetrabit*. β . H. F. 226.

White flowered Hemp-leaved Dead Nettle.

*In Hedges between Plomford and Badgen Downs —
not common. August. A.*

LAMIUM cannabinum folio vulgare. R. S. 240.

GALEOPSIS *Tetrabit*. H. F. 226.

Nettle Hemp, or Hemp-leaved Dead Nettle.

In Byfing Wood — not common. August. A.

LAMIUM cannabinum folio, flore amplo luteo, labio
purpureo. R. S. 241.

GALEOPSIS *Tetrabit* δ . H. F. 226.

Hemp-leaved Dead Nettle with a parti-coloured
Flower.

*In Mr. Jacob's new Wood-hedge at Nackington —
very uncommon. August. A.*

LAMIUM folio caulem ambiente majus et minus.
R. S. 240.

LAMIUM *amplexicaule*. H. F. 225.

Great Henbit.

In Hop-grounds — very common. June. A.

LAMIUM luteum. R. S. 242.

GALEOPSIS *Galeobdolon*. H. F. 226.

Yellow Archangel, Dead Nettle, or yellow Nettle Hemp.

In Woods and under shady Hedges — very common.

May. P.

LAMPSANA. R. S. 173.

LAPSANA *communis*. H. F. 303.

Nipple-wort.

In moist Cornfields, and by Way sides — very common.

July. A.

LAPATHUM *acetosum repens lanceolatum*.

R. S. 143.

RUMEX *Acetofella*. H. F. 136.

Sheep's Sorrel.

On dry sandy or gravelly Ground — very common.

July. P.

LAPATHUM *acetosum vulgare*. R. S. 143.

RUMEX *Acetosa*. H. F. 136.

Common Sorrel.

In Meadows and Pastures — very common. June. B.

LAPATHUM *acutum*. R. S. 142.

RUMEX *acutus*. H. F. 134.

Sharp-pointed Dock.

In uncultivated Places — very common. June. P.

LAPA-

LAPATHUM acutum minimum. R. S. 141.

An RUMICIS *acuti* varietas. H. F. 134.

The lesser sharp-pointed Dock.

In Woods — not uncommon. June. P.

LAPATHUM folio acuto crispo. R. S. 141.

RUMEX *crispus*. H. F. 134.

Sharp-pointed Dock, with curled Leaves.

On the Mud Walls beyond the Sluice — common.

July. P.

LAPATHUM folio acuto rubente. R. S. 142.

RUMEX *sanguineus*. H. F. 133.

Blood-wort.

In Davington Brooks — uncommon. July. B.

LAPATHUM maximum aquaticum sive Hydro-
lapathum. R. S. 140.

RUMEX *Britannica*. H. F. 135.

Great Water Dock.

In Ditches and boggy Places — very common. August. P.

LAPATHUM pulchrum Bononiense sinuatum.

R. S. 142.

RUMEX *pulcher*. H. F. 134.

Fiddle-Dock.

By the sides of Foot-ways about the Town — not un-
common. June. P.

LAPA-

LAPATHUM viride. R. S. 141.

An RUMICIS *acuti* varietas. H. F. 134.

The greener leaved Dock.

In shady Woods and moist Ground — not uncommon.

June. P.

LAPATHUM vulgare folio obtuso. R. S. 141.

RUMEX *obtusifolius*. H. F. 134.

Common broad-leaved Dock.

In moist Places, and by Road sides — very common.

July. P.

LAPPA major Arctium Dioscoridis. R. S. 197.

ARCTIUM *Lappa*. H. F. 304.

Burdock, or Clot burr.

Under Hedges, and by Road-sides — very common.

August. B.

LAPPA major capitulo glabro maximo. R. S. 196.

ARCTIUM *Lappa*. β . H. F. 304.

Greater Burdock, with a larger Flower.

By Road-sides, and under Hedges — common.

August. B.

LAPPA major capitulis parvis glabris. R. S. 197.

ARCTIUM *Lappa*. H. F. 304.

Burdock with small Heads.

By Road sides in Hern-hill — not uncommon. Au-

gust. B.

LAPPA

LAPPA major montana capitulis tomentosis.

ARCTIUM *Lappa*. H. F. 304.

Greater Burdock with woolly Heads.

In the Way-sides near Luddenham Court—uncommon.

August. B.

LATHYRUS luteus sylvestris dumetorum. R. S.

320.

LATHYRUS *pratensis*. H. F. 277.

Tare everlasting, or common yellow bastard Vetchling.

In Woods and Hedges — common. August. A.

LATHYRUS major latifolius. R. S. 319.

LATHYRUS *latifolius*. H. F. 276.

Broad-leaved everlasting Pea.

In a Field Hedge at Copton, and by the Road-side near Boughton Street—uncommon. July. P.

LATHYRI majoris species flore rubente et albido.
R. S. 319.

LATHYRUS *sylvestris*. H. F. 276.

Narrow-leaved everlasting Pea.

In a Hedge of the Beacon Farm—uncommon. August. P.

LAUREOLA. R. S. 465.

DAPHNE *Laureola*.

Dwarf Laurel, or Spurge-Laurel.

In Woods and under Hedges — common. March. P.

LENS

LENS palustris. R. S. 129.

LEMNA minor. H. F. 345.

Duck's Meat.

In Ponds and standing Waters — very common.
June. P.

LENTICULA aquatica trifulca. R. S. 129.

LEMNA trifulca. H. F. 344.

Ivy-leaved Duck's Meat.

In the Ditches of Ham Marsh—common. June. P.

LENTICULA palustris major. R. S. 129.

LEMNA polyrbiza. H. F. 345.

Greater Duck's Meat.

In a Ditch near Waterham—uncommon. July. A.

LEPIDIUM latifolium. R. S. 304.

LEPIDIUM latifolium. H. F. 244.

Dittander, or Pepperwort.

Near the King's Head Key in the Town—very uncommon. July. P.

LEUCANTHEMUM vulgare. R. S. 184.

CHRYSANTHEMUM *Leucanthemum*. H. F.

321.

The greater Daisie, or Ox-eye.

In Pastures, and amongst Corn—common. May. P.

LEUCO-

LEUCOJUM luteum vulgo Cheiri flore simplici.

R. S. 291.

CHEIRANTHUS *Cheiri*. H. F. 250.

Wall-flower, or wild Chier.

On old Walls — very uncommon. May. P.

* LEUCOJUM bulbosum trifolium minus.

Bauhini Pinacis. 56.

GALANTHUS *nivalis*. Lin. Syft. Natur.

H. p. 234.

The Snow Drop.

In Pastures at South-street, and at Davington —
very common. February. P.

LIGUSTRUM. R. S. 465.

LIGUSTRUM *vulgare*. H. F. 3.

Privet, or Prim.

In Woods and Hedges — very common. May. S.

LILLIUM convallium. R. S. 264.

CONVALLARIA *majalis*. H. F. 126.

Lilly Convally, or May Lilly.

In the Blean Woods — plentifully. May. P.

LIMNOPEUCE. R. S. 136.

HIPPURIS *vulgaris*. H. F. 1.

Mare's Tail.

In the Powder-Mill Waters — uncommon. May. P.

LIMO-

LIMONIUM. R. S. 200.

STATICE *Limonium*. H. F. 114.

Sea Lavender.

In the Salt Marshes—very common. August. P.

LINARIA *Elatine dicta folio acuminato.*

R. S. * 282.

ANTIRRHINUM *Elatine*. H. F. 237.

Sharp-pointed Fluellin.

In Corn Fields—common. September. A.

LINARIA *Elatine dicta folio subrotundo.*

R. S. * 282.

ANTIRRHINUM *spurium*. H. F. 237.

Round leaved Fluellin.

In the Corn fields of Preston—not uncommon. August. A.

LINARIA *lutea vulgaris*. R. S. * 281.

ANTIRRHINUM *Linaria*. H. F. 238.

Common yellow Toad-flax.

On the sides of Fields and Roads—very common. July. P.

LINUM *sylvestre angustifolium floribus dilute purpurascens vel carneis*. R. S. 362.

LINUM *tenuifolium*. H. F. 116.

Narrow-leaved Wild Flax.

Upon Beacon Hill—not common. June. P.

LINUM

LINUM sylvestre cæruleum perenne erectius flore
et capitulo majore. R. S. 362.

LINUM *perenne*. H. F. 115.

Perennial Blue Flax.

In the Isle of Shepey — *uncommon*. June. P.

LINUM sylvestre catharticum. R. S. 362.

LINUM *catharticum*. H. F. 116.

Purging Flax, Wild Dwarf Flax, Mill-mountain
On sandy and chalky Soils — *very common*. June. A.

LITHOSPERMUM, seu Miliun Solis. R. S. 228.

LITHOSPERMUM *officinale*. H. F. 65.

Gromwell, Gromil, or Graymill.

In Badgen Wood, and by Road sides — *not uncommon*.
June. P.

LONCHITIS aspera. R. S. 118.

OSMUNDA *Spicant*. H. F. 382.

Rough Spleen-wort.

In the Woods near Dunkirk — *uncommon*. July. P.

LOTI corniculatæ major species. R. S. 334.

LOTUS *corniculata*. H. F. 288. γ.

The greater Bird's-foot Trefoil.

Under Hedges near Byssing Wood — *not common*.

July. P.

G

LOTUS

LOTUS corniculata glabra minor. R. S. 334.

LOTUS *corniculata*. H. F. 288.

Bird's-foot Trefoil.

Under Hedges, and in Fields in Ospringe——common.

July. P.

LUPULUS mas et foemina. R. S. 137.

HUMULUS *Lupulus*. H. F. 369.

Hops, the male and female.

In Hedges in Preston — not uncommon. July. P.

LUTEOLA. R. S. 366.

RESEDA *Luteola*. H. F. 181.

Wild Woad, Yellow Weed, or Dyers Weed.

In Davington Church-Yard, and on barren Lands —
not uncommon — June. A.

LYCHNIS major noctiflora Dubrensis perennis.

R. S. 340.

CUCUBALUS *viscosus*. H. F. 163.

Dover Campion.

On the Cliffs between St. Margaret's and Dover —
plentifully. July. P.

LYCHNIS maritima repens. R. S. 337.

SILENE *amæna*.? H. F. 164.

Sea Campion.

In the Isle of Shepey. — uncommon. August. P.

LYCH-

LYCHNIS plumaria sylvestris simplex. R.S. 338.

LYCHNIS *Flos cuculi*. H. F. 174.

Meadow Pink, Wild Williams, Cuckow Flower.

In moist Pastures — very common. June. P.

LYCHNIS Saponaria dicta. R. S. 339.

SAPONARIA *officinalis*. H. F. 160.

Common Soap-wort.

In Hedges near Sindal and Keatings, and beyond
Sheldwich Church — uncommon. July. P.

LYCHNIS fegetum major. R. S. 338.

AGROSTEMMA *Gitbaco*. H. F. 173.

Cockle.

In Corn Fields — common. June. A.

LYCHNIS supina maritima *Ericæ facie*. R.S. 338.

FRANKENIA *lævis*. H. F. 119.

Smooth Sea-Heath.

Near Minster in Shepey — uncommon. August. P.

LYCHNIS sylvestris angustifolia caliculis turgidis
striatis. R. S. 341.

SILENE *conoidea*. H. F. 165.

Narrow-leaved Campion.

On the Sandbills near Deal — uncommon. July.

LYCHNIS sylvestris flore albo. R. S. 339.

LYCHNIS dioica. H. F. 174.

Wild white Campion.

In Woods and Hedges — not uncommon. June. P.

LYCHNIS sylvestris flore rubello. R. S. 339.

LYCHNIS dioica. H. F. 174. β .

Red flowered wild Campion.

In Woods and Hedges — common. June. P.

LYCHNIS sylvestris, quæ Behen album vulgo.

R. S. 337.

CUCUBALUS Behen. H. F. 163.

Spatling Poppy, Bladder Campion, or white
Corn Campion.

In dry Pastures, and in Corn fields — common. July. P.

LYCOPUS palustris glaber. R. S. 236.

LYCOPUS europæus. H. F. 8.

Water Horehound.

On the Banks of the River, and in boggy Places —
common. July. P.

LYSIMACHIA campestris. R. S. 311.

EPILOBIUM montanum. H. F. 141.

The greater smooth-leaved Willow-herb, or
Loose-strife.

By the sides of Ditches, and under moist Hedges —
common. July. B. LYSI-

LYSIMACHIA lutea. R. S. 282.

LYSIMACHIA vulgaris. H. F. 72.

Yellow Willow-herb, or Loose-strife.

On the swampy Ground near Stone Bridge — very uncommon. August. P.

LYSIMACHIA filiquosa glabra media five minor.

R. S. 311.

EPILOBIUM tetragonum. H. F. 141.

Middle smooth-leaved Willow-herb, or Loose-strife.

In moist Ditches, and near the River — common.

July. P.

LYSIMACHIA filiquosa glabra minor angustifolia. R. S. 311.

EPILOBIUM palustre. H. F. 141.

The least smooth coddled Willow-herb, or Loose-strife.

In moist Places — common. July. P.

LYSIMACHIA filiquosa hirsuta magno flore.

R. S. 311.

EPILOBIUM ramosum. H. F. 141.

Great flowered Willow-herb, called Codlings and Cream.

On the sides of the shooting Meadow Stream — common.

July. P.

LYSIMACHIA filiquosa hirsuta parvo flore.

R. S. 311.

EPILOBIUM hirsutum. H. F. 140.

The lesser hairy coddled Willow-herb, or Loose-strife.

On the sides of the shooting Meadow Stream — common.

July. P.

LYSI-

LYSIMACHIA speciosa, quibusdam Onagra dicta. R. S. 310.

EPILOBIUM angustifolium. H. F. 140.
Rosebay Willow-herb.

In a Wood Hedge beyond the five Mile Stone to Charing
— very uncommon. July. P.

M.

MALUS sylvestris. R. S. 452.

PYRUS Malus. H. F. 189.

The Crab Tree.

In Hedges — not uncommon. May. S.

MALVA sylvestris minor. R. S. 251.

MALVA rotundifolia. H. F. 268.

Small wild or Dwarf Mallow.

On dry Banks by Road sides — common. July to
October. A.

MALVA vulgaris. R. S. 251.

MALVA sylvestris. H. F. 268.

Common Mallow.

Under Hedges, by Way sides, and on Rubbish — very
common. July. B.

MARRUBIUM album. R. S. 239.

MARRUBIUM vulgare. H. F. 228.

White Horehound.

On Sheldwich Lees — not common. July. P.

MATRI-

MATRICARIA. R. S. 187.

MATRICARIA *Parthenium*. H. F. 321.

Feverfew.

Under Hedges, and on old Walls—common. June. P.

MELAMPYRUM *sylvaticum* flore luteo. R. S.

* 286.

MELAMPYRUM *sylvaticum*. H. F. 236.

Common Cow-wheat with a yellow Flower.

In Byfing Wood—very common. June to Auguft. A.

MELILOTUS *vulgaris*. R. S. 331.

TRIFOLIUM *Melilotus officinalis*. H. F. 282.

Common Melilot.

Amongft the Corn at Halfto—plentifully. July. B.

* MELISSA *officinalis*. Miller Botan. Off. 290.

MELISSA *officinalis*. Lin. System. Nat. II.

p. 400.

Common Garden Balm.

Under Hedges in Boughton and Davington — un-
common. July. P.

MENIANTHES *palustre triphyllum latifolium*
et angustifolium. R. S. 285.

MENYANTHES *trifolia*. H. F. 71.

Marsh Trefoil, or Buckbean.

In the moist Meadows of the Abbey—common. July. P.

MENTHA

MENTHA aquatica seu Sysimbrium. R. S. 233.

MENTHA aquatica. H. F. 223.

Water-mint.

In moist Places, and by the River side — very common.

August. P.

MENTHA seu Calamintha aquatica. R. S. 232.

MENTHA arvensis. H. F. 223.

Water-mint with whirled coronets.

On the sides of Ditches, and in Corn fields — common.

August. P.

MENTHA spicata angustifolia glabra spica latiore.

R. S. 234.

MENTHA spicata. H. F. 221.

Spear-mint.

*On the Side of the River opposite the Powder Stove —
very uncommon.* August. P.

MENTHASTRUM folio rugoso rotundiore spontaneum flore spicato odore gravi. R. S. 234.

MENTHA rotundifolia. H. F. 221.

Round leaved Horse-mint.

In a Meadow behind King's Mill — uncommon. August. P.

MENTHASTRUM spicatum folio longiore candidante. R. S. 234.

MENTHA longifolia. H. F. 221.

Long leaved Horse-mint.

*On the Banks of the River above Ospringe Church —
not uncommon.* August. P. MER-

MERCURIALIS annua glabra vulgaris, Mercurialis mas et Foemina. R. S. 139.

MERCURIALIS annua. H. F. 371.

French Mercury.

On Rubbish and in Gardens — not common. September. A.

MERCURIALIS perennis repens. Cynocrambe dicta. R. S. 138.

MERCURIALIS perennis. H. F. 371.

Dog's Mercury.

In Woods and under Hedges — very common. April. P.

MESPILUS Apii folio sylvestris spinosa, five Oxyacantha. R. S. 453.

CRATÆGUS Oxyacantha. H. F. 188.

The white Thorn, or Hawthorn, called Quick.

In Woods and Hedges — very common. May. S.

MESPILUS Apii folio sylvestris non spinosa. R. S. 453.

CRATÆGUS torminalis. H. F. 188.

The common or wild Service-tree, or Sorb.

In the Woods about Chart — uncommon. April or May. S.

MILLEFOLIUM vulgare. R. S. 183.

ACHILLEA Millefolium. H. F. 324.

Common Yarrow, Millfoil, or Nose-bleed.

In Meadows and Pastures — very common. June. P.

MOL.

MOLLUGO montana minor Gallio albo fimilis.

R. S. 224.

GALIUM montanum. H. F. 56.

Small mountain bastard Madder, or mountain
Ladies Bedstraw.

In the Corn fields about Sindal—not uncommon. July. P.

MOLLUGO vulgator. R. S. 223.

GALIUM Mollugo. H. F. 56.

Wild Madder, or great bastard Madder.

In Hedges and bushy Places—very common. July. P.

MOLLUGINIS vulgatoris varietas minor.

R. S. 224.

GALIUM palustre. H. F. 57.

White Ladies Bedstraw.

On the sides of the shooting Meadow Stream—common.

July. P.

MOSCHATELLINA foliis fumarizæ bulbosæ.

R. S. 267.

ADOXA Moschatellina. H. F. 150.

Tuberous Moschatell.

In Woods and Hedges—very common. April. P.

MYOSOTIS scorpioides hirsuta. R. S. 229.

MYOSOTIS scorpioides. H. F. 65.

Moufe-ear Scorpion-grafs.

In Woods, on old Walls and dry Pastures—common.

April to August. P. MYOSO-

MYOSOTIS scorpioides palustris. R. S. 229.

MYOSOTIS scorpioides. H. F. 65. 8.

Water Scorpion-grass.

On the sides of Ore Mill-pond—not common. April
to August. P.

MYRRHIS sylvestris feminibus asperis. R. S. 220.

SCANDIX Anthriscus. H. F. 108.

Small Hemlock - Chervil with rough Seeds.

Under Hedges, and by the sides of Ditches — common.
May. A.

N.

NAPUS sylvestris. R. S. 295.

BRASSICA Napus. H. F. 253.

Wild Navew.

Amongst Corn, and by the sides of Ditches — common.
May. B.

NARCISSUS sylvestris pallidus calyce luteo.
R. S. 371.

NARCISSUS Pseudo Narcissus. H. F. 123.

Wild English Daffodil.

In the Orchards near Selling-Street—very uncommon.
April. P.

NASTURTIUM fupinum capsulis verrucosis, five

Coronopus Ruellii. R. S. 304.

COCHLEARIA Coronopus. H. F. 248.

Swine's Cress.

On the sides of Roads and Foot-paths — very common.
July. A.

NEPETA

NEPETA major vulgaris. R. S. 237.

NEPETA Cataria. H. F. 220.

Nep, or Cat-mint.

Under the West Hedge of King's-Field — not common.

July. P.

NUMMULARIA. R. S. 283.

LYSIMACHIA Nummularia. H. F. 73.

Money-wort, or Herb Two-pence.

In moist Hedges at Luddenham, and near Provender

— not common. June. P.

NUMMULARIA minor flore purpurascente.

R. S. 283.

LYSIMACHIA tenella. H. F. 73.

Purple flowered Money-wort.

In moist Meadows by the Abbey — not uncommon.

July. P.

NYMPHÆA alba. R. S. 368.

NYMPHÆA alba. H. F. 206.

White Water Lilly.

In Ham Ponds near Sandwich — uncommon. July. P.

NYMPHÆA lutea. H. F. 368.

NYMPHÆA lutea. H. F. 206.

Yellow Water Lilly.

In Ore Mill Pond — uncommon. July. P.

OENAN-

O.

OENANTHE *aquatica*. R. S. 210.OENANTHE *fistulosa*. H. F. 104.

Water Dropwort.

*In the Ditches near Fairbrook—common. July. P.*OENANTHE *Cicutæ facie*. Lobellii. R. S. 210.OENANTHE *erocata*. H. F. 105.

Hemlock Dropwort.

*In the Water-lane between Sitingbourn and Milton—
very uncommon. June. P.*

ONOBRYCHIS seu Caput gallinaceum. R. S. 327.

HEDYSARUM *Onobrychis*. H. F. 281.

Saintfoin or Cock's-head.

On the chalky Hills — common. June. P.

OPHIOGLOSSUM. R. S. 128.

OPHIOGLOSSUM *vulgatum*. H. F. 382.

Adders Tongue.

*In Meadows at Ospringe and Luddenham — not un-
common. May. P.*OPULUS *Ruellii*. R. S. 460.VIBURNUM *Opulus*. H. F. 112.

Water Elder, or Cherrywood.

*In Badgen Wood, and in Hedges — not uncommon.
June. S.*

H

ORCHIS

ORCHIS abortiva rufa sive Nidus avis. R. S. 382.

OPHRYS *Nidus avis*. H. F. 338.

Birds nest Orchis.

In Byfing and Jud's Woods — *very uncommon*.

June. P.

ORCHIS alba bifolia minor calcari oblongo.

R. S. 380.

ORCHIS *bifolia*. H. F. 333.

Butterfly Orchis.

In Badgen and Cockset Woods — *not common*.

May. P.

ORCHIS anthropophora oreades. R. S. 397.

OPHRYS *anthropophora*. H. F. 340.

The Green Man Orchis.

On chalky Banks — *common*. June. P.

ORCHIS fuciflora galea et alis purpurascens.

R. S. 379.

OPHRYS *apifera*. H. F. 340.

The Bee Orchis.


In Meadows in Ospringe — *not uncommon*. June. P.

ORCHIS galea et alis fere cinereis. R. S. 378.

ORCHIS *militaris*. H. F. 335.

The Man Orchis.

On chalky bushy Banks beyond Whitehill, Ospringe —
not common. June. P.



C. Hall sc.
Orchis Morio femina flore pleno

ORCHIS magna latifolia galea fusca vel nigricante.

R. S. 378.

ORCHIS *purpurea*. H. F. 334.

The purple Man Orchis.

In Cockset and Jud's Woods—uncommon. May. P.

ORCHIS morio foemina. R. S. 377.

ORCHIS *morio*. H. F. 333.

Female Orchis, or Female Fools Stones.

In Meadows — very common. In a Meadow near Cades in Ospringe, some of a Rose Colour, some white, and a very few with double Flowers. May and June. P.

ORCHIS morio mas foliis maculatis. R. S. 376.

ORCHIS *mascula*. H. F. 333.

Male Orchis, or Male Fools Stones.

In Hedges and Meadows—very common. May. P.

ORCHIS myodes galea et alis herbidis. R. S. 379.

OPHRYS *muscifera*. H. F. 340.

The Fly Orchis.

Among the Bushes in Ospringe Parsonage Meadows— not uncommon. June. P.

ORCHIS odorata moschata. R. S. 378.

OPHRYS *Monorchis*. H. F. 339.

Yellow or Musk Orchis.

On the chalky Banks of Ospringe Parsonage Meadows — uncommon. June. P.

ORCHIS palmata minor flore luteo-viridi.

R. S. 381.

SATYRIUM viride. H. F. 337.

Frog Satyrion or Orchis.

On dry grassy Banks near Whitehill ——— uncommon.

June. P.

ORCHIS palmata pratensis latifolia longis calcaribus. R. S. 380.

ORCHIS latifolia. H. F. 335.

Male handed Orchis, or Satyrion royal.

On moist boggy Grounds about Ore Mill ——— not uncommon. June. P.

ORCHIS palmata speciosiore thyrsifolia maculato.

R. S. 381.

ORCHIS maculata. H. F. 335.

Female handed Orchis, or Satyrion royal.

In Cbalk Pits — common. June. P.

ORCHIS pannonica. R. S. 377.

ORCHIS ustulata. H. F. 334.

Little purple flowered Orchis.

On the South side of Chatham Hill ——— plentifully.

May. P.

ORCHIS purpurea spica congesta pyramidalis.

R. S. 377.

ORCHIS pyramidalis. H. F. 334.

Purple late flowering Orchis.

On chalky Banks in Ospringe — common. July. P.

ORCHIS *spiralis alba odorata*, five *Triorchis*.

R. S. 378.

OPHRYS *spiralis*.

Triple Ladies Traces.

In Ospringe Parsonage Meadows — *uncommon*.

September. P.

ORCHIS five *Testiculus sphegodes hirsuto flore*.

R. S. 380.

OPHRIS *apifera*. H. F. 340. β .

Humble Bee Orchis with green Wings.

On the Banks of Whitehill in Selling — *uncommon*.

April. P.

ORIGANUM *Onites*. R. S. 236.

ORIGANUM *Onites*. H. F. 229.

Pot Marjoram.

In Waldershare Park — *very uncommon*. August. P.

ORIGANUM *vulgare spontaneum*. R. S. 236.

ORIGANUM *vulgare*. H. F. 229.

Wild Marjoram.

Under Hedges, and in bushy Places — *very common*.

July. P.

ORNITHOPODIUM *radice nodosa*. R. S. 326.

ORNITHOPUS *perpusillus*. H. F. 280.

Birds Foot.

In Broom fields at Hernhill — *common*. May to Au-

gust. P.

OROBANCHE major Garyophyllum olens.

R. S. * 288.

OROBANCHE major. H. F. 232.

Broom-rape.

In Broom fields — not uncommon. June. B.

OROBUS fylvaticus foliis oblongis glabris.

R. S. 324.

OROBUS tuberosus. H. F. 274.

Wood Pea, or Heath Pea.

In Jud's Wood — very common. May. P.

OSTRYA Ulmo similis, fructu in umbilicis foliaceis.

R. S. 451.

CARPINUS. Betulus. H. F. 360.

The Horn-beam Tree, or Horse-Beech.

In Woods — common. May. S.

OXYS alba. R. S. * 281.

OXALIS Acetosella. H. F. 173.

Wood Sorrel.

In Byfing and Jud's Woods — common. June. A.

P.

PAPAYER corniculatum luteum. R. S. 309.

CHELIDONIUM Glaucium, H. F. 202.

Yellow horned Poppy.

On the Beach at Sea-Salter — common. July. A.

PAPAYER

PAPAVER corniculatum luteum. Chelidonia
dictum. R. S. 309.

CHELIDONIUM majus. H. F. 201.

The greater Celandine.

Among Rubbish, and on old Walls—common. June. P.

PAPAVER laciniato folio capitulo brevior glabro
annuum, Rhœas dictum. R. S. 308.

PAPAVER Rhœas. H. F. 203.

Red Poppy, or Corn Poppy.

In Corn fields — very common. July. A.

PARIETARIA. R. S. 158.

PARIETARIA officinalis. H. F. 376.

Pellitory of the Wall.

On old Walls — very common. July. P.

PARONYCHIA vulgaris. R. S. 292.

DRABA verna. H. F. 243.

Common Whitlow Grass.

On old Walls and dry Pastures — very common.

April. A.

PASTINACA sylvestris latifolia. R. S. 206.

PASTINACA sativa. H. F. 109.

Wild Parsnep.

*On the chalky Bank by the Roadside near Wilderton—
uncommon. July. B.*

PEDI-

PEDICULARIS seu Crista galli lutea. R. S.
* 284.

RHINANTHUS *Crista galli*. H. F. 234.

Yellow Rattle, or Cocks-comb.

In Byfing Wood, and by Road sides—common. July. A.

PEDICULARIS palustris rubra elatior. R. S.
* 284.

PEDICULARIS *palustris*. H. F. 236.

Great Marsh red Rattle, Cocks-comb, Louse-wort.

About the Powder Mills at Ore—not common. June. A.

PEDICULARIS pratensis rubra vulgaris. R. S.
* 284.

PEDICULARIS *sylvatica*. H. F. 236.

Red Rattle, Cocks-comb, Louse-wort.

In Jud's Wood—common. June. P.

PENTAPHYLLOIDES *Argentina dicta*. R. S.
256.

POTENTILLA *Argentina*. H. F. 195.

Wild Tansey, or Silver-weed.

By the sides of Roads and Paths—common. July. P.

PENTAPHYLLUM *erectum* foliis profunde
sectis subtus argenteis flore luteo. R. S. 255.

POTENTILLA *argentea*. H. F. 196.

Tormentil Cinquefoil.

In a Road hedge near Newnham.—uncommon. May. P
PENTA-

PENTAPHYLLUM vulgatissimum. R. S. 225.

POTENTILLA reptans. H. F. 197.

Common Cinquefoil, or Five-fingered Grass.

By the sides of Roads and Paths—common. June. P.

PENTAPTEROPHYLLON aquaticum, flosculis
ad foliorum nodos. R. S. 316.

MYRIOPHYLLUM verticillatum. H. F. 357.

Verticillated Water-millfoil.

In the Ditches of Graveney Marsh — not uncommon.
July. P.

PERCEPIER Anglorum. R. S. 159.

APHANES arvensis. H. F. 60.

Pursley-piert.

Upon barren Lands and old Walls—common. May. A.

PERSICARIA latifolia geniculata caulibus macu-
latis. R. S. 145.

POLYGONUM Persicaria. H. F. 147. β.

Dead or spotted Arsmart.

In Fields, and on Rubbish — common. August. A.

PERSICARIA maculosa. R. S. 145.

POLYGONUM Persicaria. H. F. 147.

Dead or spotted Arsmart.

In Meadows and Ditches—very common. Sept. A.

PERSI-

PERSICARIA mitis major foliis pallidioribus.

R. S. 145.

POLYGONUM *pensylvanicum*. H. F. 148.

Greater dead Arsmart with pale Leaves.

On a Dungbill at Staple Street — uncommon.

August. A.

PERSICARIA *salicis folio perennis*, Potamogiton

angustifolium dicta. R. S. 145.

POLYGONUM *amphibium*. H. F. 147.

Perennial Willow-leaved Arsmart.

On the sides of the Abby Ditches — not common.

July. P.

PERSICARIA *vulgaris acris*, seu Hydropiper.

R. S. 144.

POLYGONUM *Hydropiper*. H. F. 148.

Lakeweed, Arsmart, Water-pepper.

In moist Places and wet Ditches — very common.

August. A.

PETASITES. R. S. 179.

TUSSILAGO *Petasites*. H. F. 315.

Common Butter Bur, Pestilent-wort.

Near Waterham in Hernhill — uncommon. March. P.

• PETROSELINUM *vulgare*. Miller. Botan.

Officinal. 324.

APIUM *Petroselinum*. Linnæi System. Natur.

II. p. 217.

Common Parsley.

On the West Church-yard Wall — very uncommon.

June. B.

PEU-

PEUCEDANUM. R. S. 206.

PEUCEDANUM *officinale*. H. F. 101.

Hog-fennel, or Sulphur-wort.

On the Wall leading to Thorn Creek — plentifully.
June. P.

PELLANDRIUM vel Cicutaria aquatica quorundam. R. S. 215.

PELLANDRIUM *aquaticum*. H. F. 107.

Water Hemlock.

On boggy Ground between Staple-Street and Hernhill Church — uncommon. July. B.

PHYLLITIS feu Lingua cervina vulgo. R. S. 116.

ASPENIUM *Scolopendrium*. H. F. 384.

Hart's Tongue.

In shady Lanes, hollow Ways, and on Walls — common. P.

PILOSELLA repens. R. S. 170.

HIERACIUM *Pilosella*. H. F. 298.

Common creeping Mouse-ear.

On dry sandy Banks — common. May. A.

PIMPINELLA *Saxifraga*. R. S. 213.

PIMPINELLA *major*. H. F. 110.

Great Burnet-saxifrage.

In a thick Hedge beyond Whitehill — uncommon.
August. P.

PIMPI-

PIMPINELLA Saxifraga minor. R. S. 213.

PIMPINELLA Saxifraga. H. F. 111. β .

Small Burnet-saxifrage with divided Leaves.

In Meadows in Hernhill — not uncommon. August. P.

PIMPINELLA Saxifraga minor foliis sanguiforbæ. R. S. 213.

PIMPINELLA Saxifraga. H. F. 111.

The lesser round leaved Burnet-saxifrage.

In dry Pastures — common. August and September. P.

PLANTAGO aquatica. R. S. 257.

ALISMA Plantago. H. F. 137.

Great Water Plantain.

In moist Places, and on the sides of Ponds — common. August. P.

PLANTAGO foliis laciniatis, Coronopus dicta.

PLANTAGO Coronopus. H. F. 52.

Buck's-horn Plantain, Star of the Earth.

In the Abby Meadows — very common. August. A.

PLANTAGO latifolia vulgaris. R. S. 314.

PLANTAGO major. H. F. 51.

Great Plantain, Way-bread.

By Way sides — very common. July. P. The

Plantago panicula sparsa, found in Perry field, is an elegant variety of this Plant, but very uncommon. PLAN-

PLANTAGO major incana. R. S. 314.

PLANTAGO media. H. F. 51.

Hoary Plantain.

By Road sides — common. July. P.

PLANTAGO marina. R. S. 315.

PLANTAGO Loeftingii. H. F. 52.

Sea Plantain.

On the sides of the Ditches and Sea Walls — common.

July. P.

PLANTAGO maritima. H. F. 52.

Narrow leaved Plantain.

In the Brent Marshes — not uncommon. July. P.

PLANTAGO quinquenervia. R. S. 314.

PLANTAGO lanceolata. H. F. 52.

Rib-wort, or Rib wort Plantain.

In Fields and Meadows — very common. June to

August. P.

POLYGALA. R. S. * 287.

POLYGALA vulgaris. H. F. 271.

Milk-wort.

In dry Fields, and under Hedges — very common.

June. P.

POLYGONUM *brevi angustoque folio.* R. S. 146.

POLYGONUM *aviculare.* H. F. 149. β .

Knot-grass with a small narrow Leaf.

By Road sides—common. June to September. A.

POLYGONUM *mas vulgare.* R. S. 146.

POLYGONUM *aviculare.* H. F. 149.

Common Knot-grass.

In uncultivated Ground, and by Way sides — very common. June to October. A.

POLYGONUM *oblongo angustoque folio.*

R. S. 146.

POLYGONUM *aviculare.* H. F. 149. γ .

Knot-grass with a more oblong narrow Leaf.

In Corn fields at Hernhill — common. June to September. A.

POLYPODIUM. R. S. 117.

POLYPODIUM *vulgare.* H. F. 387.

Common Polypody.

In Byfing Wood, and on Banks of hollow Ways — common. P.

POPULAGO; five *Caltha palustris.* R. S. 272.

CALTHA *palustris.* H. F. 214.

Marsh Marigold.

In the Brent Marshes—very common. April. P.

POPU-

POPULUS alba. R. S. 446.

POPULUS alba. H. F. 370.

The White Poplar, or Abele Tree.

In moist Woods, and boggy Places — common.

March. S.

POPULUS Lybica Plinii. R. S. 446.

POPULUS tremula. H. F. 370.

The Asp, or Trembling Poplar.

In moist Places in Hernhill — not uncommon.

March. S.

POPULUS major. R. S. 446.

POPULUS nigra. H. F. 370.

The Black Poplar.

In moist boggy Places in Hernhill — not uncommon.

March. S.

PORTULA. R. S. 368.

PEPLIS Portulaca. H. F. 128.

Water Purslane.

In the Brent Meadows — not uncommon. September. P.

POTAMOGETON foliis pennatis. R. S. 150.

MYRIOPHYLLUM spicatum. H. F. 357.

Feathered or spiked Water Millfoil.

In the Powder Mill Waters, and Graveney Marsh

Dykes — common. July. P.

POTAMOGITON seu Fontalis crispa. Tribulus aquaticus. R. S. 149.

POTAMOGETON *crispum*. H. F. 61.

The greater Water Caltrops.

In the Powder Mill Waters—common. June. P.

POTAMOGETON seu Fontalis media lucens.
R. S. 149.

POTAMOGETON *ferratum*. H. F. 61.

The lesser Water Caltrops, or Frog's Lettuce.

In the River near the Powder Stove—not uncommon.
June. P.

POTAMOGITON gramineum latiusculum foliis
et ramificationibus dense stipatis. R. S. 149.

POTAMOGETON *gramineum*. H. F. 62.

Grass leaved Pond-weed.

In the shallow Waters about the Powder Mills — not
common. P.

POTAMOGITON maritimum ramosissimum grandiusculis capitulis, capillaceo folio nostras.
R. S. 150.

POTAMOGETON *marinum*. H. F. 63.

Sea Pond-weed.

In Dykes near Sheerness — uncommon. August. P.

POTAMOGITON millefolium, seu foliis gramineis
ramosum. R. S. 150.

POTAMOGETON *peelinatum*. H. F. 62.

Fine or Fennel-leaved Pond-weed.

In Ospringe River — not common. June. P.

POTAMOGETON perfoliatum. R. S. 149.

POTAMOGETON perfoliatum. H. F. 60.

Perfoliated Pond-weed.

In Ospringe River — common. July. P.

POTAMOGETON rotundifolium. R. S. 148.

POTAMOGETON natans. H. F. 60.

Broad leaved Pond-weed.

In a Pond near Uples — uncommon. August. P.

PRIMULA pratensis inodora lutea. R. S. 284.

PRIMULA veris. H. F. 70. β .

Great Cowslips, or Oxslips.

In thick Hedges beyond Whitehill — not common.

April. P.

PRIMULA veris major. R. S. 284.

PRIMULA veris. H. F. 71. α .

Common Pagils, or Cowslips.

In Meadows and Hedges — very common. April. P.

PRIMULA veris vulgaris. R. S. 284.

PRIMULA vulgaris. H. F. 70.

Common Primrose.

In Woods and Hedges — very common. February to
May. P.

PRUNELLA. R. S. 238.

PRUNELLA vulgaris. H. F. 231.

Common Self-heal.

In Meadows — common. August. P.

PRUNUS sylvestris. R. S. 48.

PRUNUS spinosa. H. F. 186.

The Black Thorn, or Sloe Tree.

In Hedges — very common. April. S.

PRUNUS sylvestris major. R. S. 462.

PRUNUS infititia. H. F. 186.

The black Bullace Tree.

In some Hedges near Rickesey-Lane in Selling —
uncommon. April. S.

PRUNUS sylvestris fructu majore albo. R. S. 462.

PRUNUS infititia. H. F. 186. γ.

The white Bullace Tree.

In Hedges at Cades in Ospringe, and at Selling —
uncommon. April. S.

PRUNUS sylvestris fructu rubro acerbo et ingrato.

R. S. 463.

PRUNUS infititia. H. F. 186. β.

The Scad Tree.

In Hedges — not uncommon. April. S.

PTARMICA. R. S. 183.

ACHILLEA *Ptarmica*. H. F. 325.

Sneeze-wort, Bastard Pellitory, or Goose-tongue.

In Davington Brooks — *uncommon*. August. P.

PULEGIUM. R. S. 235.

MENTHA *Pulegium*. H. F. 224.

Penny-royal, or Pudding-grafs.

In a small Pond at Elvyland in Ospringe — *uncommon*.

PYRASTER seu *Pyrus sylvestris*. R. S. 452.

PYRUS *communis*. H. F. 189.

The Wild Pear Tree.

In the Wood leading from Keneways to Wilderton —
very uncommon. April. S.

Q.

QUERCUS *latifolia*. R. S. 440.

QUERCUS *Robur*. H. F. 359.

The common Oak Tree.

In Woods — *very common*. April. P.

QUERCUS *latifolia* mas quæ brevè pediculo est,
R. S. 440.

QUERCUS *Robur*. H. F. 359. β .

The Oak Tree.

In Woods and Hedges — *common*. April. S.

RADI.

R.

RADICULA sylvestris seu palustris. R. S. 301.

SISYMBRIUM amphibium. H. F. 258. α.

Water Radish.

About the Powder-Mill Waters — not common.

June. P.

RADIOLA vulgaris serpyllifolia Millegrana'
minima. R. S. 345.

LINUM Radiola. H. F. 117.

The least Rupture-wort, or All-seed.

*On sandy Banks between Ore and Luddenham —
common. A.*

RANUNCULO five Polyanthemo aquatili albo
affine. R. S. 250.

RANUNCULUS aquatilis. H. F. 213. γ.

Fennel-leaved Crow-foot.

In Running Waters — common. May. P.

RANUNCULUS aquaticus albus, circinatis
tenuissime divisis foliis, floribus ex alis longis
pediculis innixis. R. S. 249.

RANUNCULUS aquatilis. H. F. 213. δ.

Finest leaved Water Crow-foot.

In the Brent Dykes — common. April to June. P.

RANUN.

RANUNCULUS aquatilis. R. S. 249.

RANUNCULUS aquatilis. H. F. 213.

Various leaved Water Crow-foot.

In the shooting Meadow Stream — common.

May. P.

RANUNCULUS aquatilis hederaceus albus.

R. S. 249.

RANUNCULUS hederaceus. H. F. 212.

Ivy leaved Water Crow-foot.

By the Wall that leads to Thorn — common. May. P.

RANUNCULUS aquatilis omnino tenuifolius.

R. S. 249.

RANUNCULUS aquatilis. H. F. 213. β .

Fine leaved Water Crow-foot.

In Cspringe River near Whitehill — common. April
to June. P.

RANUNCULUS arvorum. R. S. 248.

RANUNCULUS arvensis. H. F. 212.

Corn Crow-foot.

In Corn fields — very common. June. A.

RANUNCULUS bulbosus. R. S. 247.

RANUNCULUS bulbosus. H. F. 211.

Round rooted or bulbous Crow-foot, or Butter-
cups

In Meadows — very common. May. P.

RANUN-

RANUNCULUS flammeus major. R. S. 250.

RANUNCULUS *Lingua*. H. F. 210.

Great Spear-wort.

In the old Haven near Sandwich — very uncommon.

May. P.

RANUNCULUS flammeus minor. R. S. 250.

RANUNCULUS *Flammula*. H. F. 210.

The lesser Spear-wort.

In the shooting Meadow Stream — common. June to
September. P.

RANUNCULUS nemorosus dulcis secundus

Tragi. R. S. 248.

RANUNCULUS *Auricomus*. H. F. 211.

Sweet wood Crow-foot, or Goldilocks.

In Woods and under Hedges — common. April. P.

RANUNCULUS palustris. R. S. 249.

RANUNCULUS *sceleratus*. H. F. 212.

Round leaved Water Crow-foot.

On the Banks of the Brent Marshes — common.
June. A.

RANUNCULUS pratensis erectus acris. R. S.
248.

RANUNCULUS *acris*. H. F. 211.

Upright Meadow Crow foot.

In Meadows — very common. June. P.

RANUNCULUS pratensis repens. R. S. 247.

RANUNCULUS *repens*. H. F. 210.

Common creeping Crow-foot.

On Beacon Hill — common. May. P.

RAPA

RAPA fativa rotunda. R. S. 294.

BRASSICA *Rapa*. H. F. 253.

Turneps.

In Corn fields, and by Way sides — not uncommon.

April. B.

RAPISTRUM arvorum. R. S. 295.

SINAPIS *arvensis*. H. F. 260.

Charlock, or Wild Mustard.

In Fields among Corn — common. May. A.

RAPISTRUM flore luteo, filiqua glabra articulata.

R. S. 296.

RHAPHANUS *Rhaphanistrum*. H. F. 252. β .

Yellow flowered Charlock with a jointed Cod.

In Fields among Corn — common. July. A.

RAPUNCULUS *Scabiosæ capitulo cæruleo*.

R. S. 278.

JASIONE *montana*. H. F. 329.

Hairy Sheep's Scabious.

By the Way sides near Dunkirk — uncommon.

August. A.

RESEDA vulgaris. R. S. 366.

RESEDA *lutea*. H. F. 181.

Base Rocket.

In Boughton Chalk Pits — not common. July. A.

RHAM.

RHAMNOIDES fructifera, foliis Salicis, baccis
leviter flavescentibus. R. S. 445.

HIPPOPHAE *Rhamnoides*. H. F. 368.

Sea Buck Thorn, or Sallow Thorn.

In Shepey, uncommon; near Sandown Castle —
plentifully. April. S.

RHAMNUS catharticus. R. S. 466.

RHAMNUS *catharticus*. H. F. 83.

Buck Thorn, or Purging Thorn.

In Perry Wood and Badgen Down Hedges — not
common. May. S.

RHAPHANISTRUM filiqua articulata glabra
majore et minore. R. S. 296.

RHAPHANUS *Rhaphanistrum*. H. F. 252.

White flowered Charlock with a jointed Cod.

In Fields among Corn — common. June. A.

RHAPHANUS rusticanus. R. S. 301.

COCHLEARIA *Armoracia*. H. F. 248.

Horse Radish.

On the Banks of the River, above the Sluice, and of
Ore Stream — uncommon. June. P.

RIBES nigrum vulgo dictum folio olente. R. S. 456.

RIBES *nigrum*. H. F. 85.

Black Currants, or Squinancy Berries.

In the boggy Ofier Ground near Stone Bridge — un-
common. May. S.

RIBES

* *RIBES vulgare fructu rubro.* R. S. 465.

RIBES rubrum. H. F. 84.

Currants.

In some thick Hedges in Ospringe — not common.

May. S.

ROSA sylvestris altera minor, flore albo nostras.

R. S. 455.

ROSA arvensis. H. F. 192.

White flowered Dog's Rose.

In Hedges — very common. June. S.

ROSA sylvestris inodora seu canina. R. S. 454.

ROSA canina. H. F. 192.

Red flowered wild Dog's Rose, or Hep Tree.

In Hedges — very common. June. S.

* *ROSA sylvestris odora.* R. S. 454.

ROSA Eglanteria. H. F. 191.

Sweet Briar.

In a Hedge near Ore Stray — very uncommon.

June. S.

ROSA pumila spinosissima, foliis Pimpinellæ

glabris flore albo. R. S. 455.

ROSA spinosissima. H. F. 191.

The Burnet Rose.

On the North side of Chatham Hill — uncommon.

June. S.

K

RUBE-

RUBEOLA arvensis repens cærulea. R. S. 225.

SHERARDIA arvensis. H. F. 54.

Little Field Madder.

In Cades new Wood, and in Broom fields in Hernhill — not uncommon. May. A.

RUBEOLA vulgaris quadrifolia lævis floribus purpurascentibus. R. S. 225.

ASPERULA Cynanchica. H. F. 55.

Squinancy Wort.

In Ospringe Chalk Pits, and on Badgen Down — July. P.

RUBUS Idæus spinosus fructu rubro. R. S. 467.

RUBUS Idæus. H. F. 193.

The Raspberry Bush.

In Stocking Wood, near Lees Court, and in Perry Wood — not common. May. S.

RUBUS major fructu nigro. R. S. 467.

RUBUS fruticosus. H. F. 193.

The Bramble, or Blackberry Bush.

In Woods and Hedges — very common. June. S.

RUBUS minor fructu cæruleo. R. S. 467.

RUBUS cæsius. H. F. 193.

The small Bramble, or Dew-berry Bush.

In Woods and Hedges — common. July. S.

RUBE.

K

RUSCUS.

RUSCUS. R. S. 262.

RUSCUS *aculeatus*. H. F. 373.

Knee Holly, or Butcher's-broom.

In Ospringe Woods and Hedges—common. April. P.

RUTA muraria. R. S. 122.

ASPLENIUM *Ruta muraria*. H. F. 386.

Wall Rue, White Maiden-hair, Tentwort.

On the Buttrices near the South Door of the Church —
very uncommon. P.

S.

SALICARIA vulgaris purpurea foliis oblongis.

R. S. 367.

LYTHRUM *Salicaria*. H. F. 179.

Purple-spiked Loosetrife, or Willow-herb.

In the swampy Ground near Stone-bridge — not un-
common. August. P.

SALICORNIA geniculata annua. R. S. 136.

SALICORNIA *europæa*. H. F. 1.

Marsh Sampire, Jointed Glass-wort, or Salt-wort.

In the Marshes as you go to Thorn — common. Au-
gust. A.

SALIX. R. S. 447.

SALIX *alba*. H. F. 366.

The common white Willow.

In moist Places about the Powder Mills—not common.

April. S.

SALIX

SALIX folio articulato splendente flexilis. R. S.
448.

SALIX amygdalina. H. F. 363.

The round eared shining Willow, or Almond-leaved Willow.

In a Hedge as you enter Sheldwich Road—uncommon.
May. S.

SALIX folio longissimo. R. S. 450.

SALIX vimenalis. H. F. 366.

The Osier.

In moist Places — common. April. S.

SALIX latifolia rotunda. R. S. 449.

SALIX caprea. H. F. 366.

The common round-leaved Sallow.

In Woods and Hedges — common. April. S.

SALIX pumila foliis utrinque candicantibus et lanuginosis. R. S. 447.

SALIX arenaria. H. F. 364.

Sand Willow.

On the Sand Downs near Deal—plentifully. June. S.

SAMBUCUS. R. S. 461.

SAMBUCUS nigra. H. F. 112.

Common Elder.

In Hedges — very common. May. S.

SAMBU-

SAMBUCUS humilis, seu Ebulus. R. S. 461.

SAMBUCUS *Ebulus*. H. F. 113.

Dwarf Elder, Wall-wort, or Dane-wort.

By the Road sides near Boughton-Street — not uncommon. July. P.

SAMOLUS Valerandi. R. S. 283.

SAMOLUS *Valerandi*. H. F. 79.

Round-leaved Water Pimpernel.

In the Brent Marshes — common. June. P.

SANGUISORBA minor R. S. 203.

POTERIUM *Sanguisorba*. H. F. 358.

Burnet.

On the chalky Hills in Ospringe — common. July. P.

SANICULA five Diapensia. R. S. 221.

SANICULA *europæa*. H. F. 96.

Sanicle.

In Woods — very common. May. P.

SAXIFRAGA aurea. R. S. 158.

CHRYSOSPLENIUM *oppositifolium*. H. F.

156.

Golden Saxifrage.

In Jud's Wood — not common. April. P.

SAXIFRAGA rotundifolia alba. R. S. 354.

SAXIFRAGA granulata. H. F. 159.

White Saxifrage, or Sen-green.

Upon Beacon Hill — uncommon. May. P.

SAXIFRAGA verna annua. R. S. 354.

SAXIFRAGA tridactylites. H. F. 159.

Rue-leaved Saxifrage, or Whitlow-grass.

On Houses and Walls — very common. April. A.

SCABIOSA major communior folio laciniato.

R. S. 191.

SCABIOSA arvensis. H. F. 50.

Common Field Scabious.

In moist Fields among Corn, and on Bank sides — common. August. P. With a proliferous Flower in Badgen Wood, uncommon.

SCABIOSA minor vulgaris. R. S. 191.

SCABIOSA Columbaria. H. F. 50.

The lesser Field Scabious.

In dry billy Places — not uncommon. July. P.

SCABIOSA radice succisa, flore globoso. R. S. 191.

SCABIOSA Succisa. H. F. 50.

Devil's Bit.

In Byfing Wood and the Abby Meadows — common.

August. P.

SCAN-

SCANDIX femine rostrato vulgaris. R. S. 207.

SCANDIX *Pecten Veneris*. H. F. 107.

Shepherd's Needle, or Venus Comb.

In Fields among Corn—very common. July. A.

SCORODONIA feu Salvia agreftis. R. S. 245.

TEUCRIUM. *Scorodonia*. H. F. 219.

Wood Sage.

In Byfing Wood—very common. July. P.

SCROPHULARIA aquatica major. R. S. * 283.

SCROPHULARIA *aquatica*. H. F. 239.

Water Betony, Water Figwort.

*On the Banks of the River, and in moist Ditches—
common.* July. P.

SCROPHULARIA major. R. S. * 283.

SCROPHULARIA *nodosa*. H. F. 239.

Common knobby-rooted Figwort.

In moist Hedges—common. July. P.

SEDUM minimum non acre flore albo. R. S.
270.

SEDUM *annuum*. H. F. 172.

Mountain Stone-crop.

On the Beach near Sandown Castle—plentifully.
August. P.

SEDUM

SEDUM minus hæmatoides. R. S. 269.

SEDUM reflexum. H. F. 170. β .

Yellow Stone-crop, or Prick-madam.

On the Abby Walls — common. July. P.

SEDUM minus luteum ramulis reflexis. R. S.

270.

SEDUM reflexum. H. F. 170.

Yellow reflexed Stone-crop.

On the Abby Walls — common. July. P.

SEDUM parvum acre flore luteo. R. S. 270.

SEDUM acre. H. F. 171.

Stone-crop, or Wall-pepper.

On Walls and Houses — not uncommon. July. P.

SEDUM sexangulare. H. F. 171.

Insipid Stone-crop.

Near Sheerness — plentifully. July. B.

SEMPERVIVUM majus. R. S. 269.

SEMPERVIVUM tectorum. H. F. 185.

House-leek, or Sengreen.

On Walls and tiling of Houses — not uncommon.

July. P.

SENECIO

SENECIO vulgaris. R. S. 178.

SENECIO *vulgaris*. H. F. 315.

Common Groundfel, or Simfon.

In Fields and under Hedges — very common.

May. A.

SERPILLUM citratum. R. S. 230.

THYMUS *Serpyllum*. H. F. 229. ε.

Lemon Thyme.

On Badgen Down — not common. July. P.

SERPILLUM vulgare. R. S. 230.

THYMUS *Serpyllum*. H. F. 229.

Mother of Thyme.

On dry billy Ground — very common. July. P.

SERRATULA. R. S. 196.

SERRATULA *tinctoria*. H. F. 304.

Saw-wort.

Near the late Decoy Ponds at Graveney — uncommon.

July. P.

SESELI pratense nostras. R. S. 216.

SESELI *Caruifolia*. H. F. 106.

Meadow Saxifrage.

*In the low Meadows in Hernhill, and in Shepey —
not uncommon. August. P.*

SIDERI-

SIDERITIS *anglica strumosa radice*. R. S. 242.

STACHYS *palustris*. H. F. 227.

Clowns Allheal.

In moist Places, and near the River — common.

August. P.

SIDERITIS *arvensis rubra*. R. S. 242.

GALEOPSIS *Ladanum*. H. F. 225.

Narrow leaved Allheal, or Iron-wort.

In Ospringe Corn fields — common. August. A.

SIDERITIS *humilis lato obtuso folio*. R. S. 242.

GLECHOMA *arvensis*. H. F. 224.

Upright Ground Ivy.

In Ore Corn fields — common. August. A.

* SINAPI *album filiqua hirsuta, semine albo vel ruffo*. R. S. 295.

SINAPIS *alba*. H. F. 260.

White Mustard.

By the Way side in the Brent Marsh — very uncommon.

August. A.

SINAPI *fativum secundum*. R. S. 295.

SINAPIS *nigra*. H. F. 260.

Common Mustard.

In Shepey amongst Corn, and by Way sides — plentifully. July. A.

SISYM.

SISYMBRIUM Cardamine seu Nasturtium aquaticum. R. S. 300.

SISYMBRIUM Nasturtium.

Water Cress.

By the River sides — very common. July. B. or P.

SISYMBRIUM hirsutum. R. S. 233.

MENTHA aquatica. H. F. 223. γ.

Water Mint.

On the sides of the River — very common. July. P.

SIUM aromaticum. Sison officinarum. R. S. 211.

SISON Amomum. H. F. 104.

Bastard Stone Parsley.

In a Hedge near Thorn — not uncommon. August. P.

SIUM arvense sive segetum. R. S. 211.

SISON segetum. H. F. 104.

Corn Parsley, or Honewort.

In moist Hedges near Ewell — not uncommon. July. B.

SIUM latifolium foliis variis. R. S. 211.

SIUM latifolium. H. F. 103.

Great Water Parsnep.

In marshy Ground, and by the River side — common. August. P.

SIUM

SIUM pusillum foliis variis. R. S. 212.

SISON inundatum. H. F. 104.

The least Water Parsnep.

In the Dykes of Nagden Marsh — not common.

June. A.

SIUM umbellatum repens. R. S. 211.

SIUM nodiflorum. H. F. 103.

Creeping Water Parsnep.

In the Abby Ditches — common. August. P.

SMYRNIUM. R. S. 208.

SMYRNIUM Olusatrum. H. F. 109.

Alexanders.

In a Hedge as you enter Graveney Marsh — uncommon.

June. B.

SOLANUM lignosum seu Dulcamara. R. S. 255.

SOLANUM Dulcamara. H. F. 78.

Woody Night-shade, or Bitter-sweet.

In moist Woods and Hedges — common. July. P.

* SOLANUM pomo spinoso oblongo flore calathoides, Stramonium vulgo dictum. R. S. 266.

DATURA Stramonium.. H. F. 78.

Thorn Apple.

On Dungbills — very uncommon. August. A.

SOLA-

SOLANUM vulgare. R. S. 265.

SOLANUM nigrum. H. F. 78.

Common Night-shade.

On Rubbish and Dungbills — very common. July. A.

SONCHUS asper laciniatus. R. S. 163.

SONCHUS oleraceus. H. F. 294. γ .

Prickly Sow-thistle with jagged Leaves.

On Rubbish and in cultivated Grounds — common.

August. A.

SONCHUS asper non laciniatus. R. S. 163.

SONCHUS oleraceus. H. F. 294. δ .

Prickly Sow-thistle with less jagged Leaves.

In cultivated Grounds, and on Rubbish — common.

June to August. A.

SONCHUS lævis. R. S. 162.

SONCHUS oleraceus. H. F. 294. α .

Common Sow-thistle.

On Rubbish, in Fields and cultivated Grounds — very common. June to September. A.

SONCHUS lævis minor paucioribus laciniis.
R. S. 163.

SONCHUS oleraceus. H. F. 294. β .

Smooth Sow-thistle with less jagged Leaves.

On Rubbish, in Fields and cultivated Grounds —

June to September. A.

L

SONCHUS

SONCHUS repens, multis Hieracium majus.
R. S. 163.

SONCHUS arvensis. H. F. 295.

Tree Sow-thistle.

In Fields and Hedges — common. August. P.

SORBUS sylvestris foliis domesticæ similis.

R. S. 452.

SORBUS aucuparia. H. F. 189.

Mountain Ash, Quicken Tree, Roan Tree.

In the Blean Woods — not uncommon. May. S.

SPARGANIUM non ramosum. R. S. 437.

SPARGANIUM erectum. H. F. 346. β .

Bur-reed not branched.

*In the Ditches of Ham and Graveney Marshes —
common. July. P.*

SPARGANIUM ramosum. R. S. 437.

SPARGANIUM erectum. H. F. 346.

Great branched Bur-reed.

In Ponds near Ewell — not uncommon. July. P.

SPERGULA maritima flore parvo cæruleo, semine
vario. R. S. 351.

ARENARIA marina. H. F. 169. γ .

Small flowered Sea-spurrey.

In Shepey Island — not uncommon. July. P.

SPER-

SPERGULA *purpurea*. R. S. 163.

ARENARIA *rubra*. H. F. 169.

Purple Spurrey, Purple flowered Chickweed.

On sandy Grounds in Hernhill — not uncommon.

July. A.

SPHONDYLUM. R. S. 205.

HERACLEUM *Sphondylium*. H. F. 102.

Cow Parsnep.

On the sides of Fields, and in Hedges — common.

July. B.

STATICE *montana minor*. R. S. 203.

STATICE *Armeria*. H. F. 114.

Thrift, or Sea Gilly-Flower.

In all the Salt Marshes — very common. July. P.

STELLARIA. R. S. 289.

CALLITRICHE *verna*. H. F. 2.

Vernal Star wort, Water wort, Star-headed Chickweed.

In the Powder Mill Streams — common. April. A.

STELLARIA *aquatica foliis longis tenuissimis*.

R. S. 290.

CALLITRICHE *autumnalis*. H. F. 2.

Autumnal Star-wort.

In the River opposite the Gunpowder Stove — not uncommon. September. A.

STRATIOTES foliis Afari, semine rotundo.

R. S. 290.

HYDROCHARIS *Morsus ranae*. H. F. 372.

Frog-bit.

In the Dykes of Ham and Graveney Marshes —
common. June. P.

SYMPHYTUM magnum. R. S. 230.

SYMPHYTUM officinale. H. F. 68.

Comfrey.

On the Banks of the Powder Mill Waters — uncommon.
July. P.

T.

TAMNUS racemosa, flore minore luteo pallescente. R. S. 262.

TAMUS communis. H. F. 369.

Black Briony.

Under Hedges, and among Bushes — common. June. P.

TANACETUM. R. S. 188.

TANACETUM vulgare. H. F. 310.

Common Tanfy.

By Way sides, near Queen Court and Uplees — very
uncommon. August. P.

TAXUS. R. S. 445.

TAXUS baccata. H. F. 372.

The Yew Tree.

In Woods and Hedges on a chalky Soil — common.
April. S. THA-

THALICTRUM seu Thaliolum majus. R. S.
203.

THALICTRUM flavum. H. F. 216.

Meadow Rue.

In the shooting Meadow Stream — uncommon.

June. P.

TITHYMALUS characias Amygdaloides. R. S.
312.

EUPHORBIA Amygdaloides. H. F. 184.

Wood Spurge.

In Woods, and under Hedges in Ospringe — very
common. April. P.

TITHYMALUS helioscopius. R. S. 313.

EUPHORBIA Helioscopia. H. F. 183.

Sun Spurge, or Wart-wort.

In Meadows, and on Dung-hills — common. July. A.

TITHYMALUS leptophyllos. R. S. 313.

EUPHORBIA exigua. H. F. 182.

Dwarf Spurge, Small annual Spurge.

In Corn fields and Gardens — common. July. A.

TITHYMALUS parvus annuus, foliis subrotundis,
non crenatis, Peplus dictus. R. S. 313.

EUPHORBIA Peplus. H. F. 182.

Petty Spurge.

In cultivated Grounds — common. July. A.

TITHYMALUS platyphyllos Fuchsii. R. S.
312.

EUPHORBIA platyphyllos. H. F. 184.
Broad leaved Spurge.

In Corn fields near Thorn Creek — common.
July. P.

TORMENTILLA. R. S. 257.

TORMENTILLA erecta. H. F. 197.
Tormentil Septfoil.

In Woods and Hedges — common. June. P.

TORMENTILLA reptans. R. S. 257.

TORMENTILLA reptans. H. F. 198.
Creeping Tormentil.

In Broom fields at Hernhill — common. July. P.

TRAGOPOGON luteum. R. S. 171.

TRAGOPOGON pratense. H. F. 293.

Yellow Goat's Beard, Go to Bed at Noon.

In Fields and Pastures — common. June. B.

TRAGOPOGON purpureum. R. S. 172.

TRAGOPOGON porrifolium. H. F. 293.

Purple Goat's Beard.

In a Marsh near Sheerness — uncommon.

June. B.

TRICHO-

TRICHOMANES. R. S. 119.

ASPLENIUM *Trichomanes*. H. F. 385.

English black Maiden-Hair.

On the Walls of the Church and Abby——not common. P.

TRIFOLIUM *arvense humile spicatum seu*

Lagopus. R. S. 330.

TRIFOLIUM *arvense*. H. F. 285.

Hare's Foot, or Hare's Foot Trefoil.

Upon Beacon Hill——not uncommon. July. A.

TRIFOLIUM *cochleatum folio cordato maculato*.

R. S. 333.

MEDICAGO *arabica*. H. F. 288.

Heart Trefoil, or Clover.

In the Abby Meadows——not uncommon. June. A.

TRIFOLIUM *cochleatum modiolis spinosis*.

R. S. 333.

MEDICAGO *arabica*. H. F. 288. γ.

Hedge-hog Trefoil, or Clover.

In Pastures near Sheerness——not common. May. A.

TRIFOLIUM *fragiferum*. R. S. 329.

TRIFOLIUM *fragiferum*. H. F. 286.

Strawberry Trefoil.

In the Abby Meadows——common. August. P.

TRIFO-

TRIFOLIUM flosculis albis, in glomerulis oblongis asperis, cauliculis proxime adnatis.

R. S. 329.

TRIFOLIUM scabrum. H. F. 285.

Oval headed Trefoil.

In Pastures in Shepey — common. June. A.

TRIFOLIUM lupulinum alterum minus. R. S.

330.

TRIFOLIUM procumbens. H. F. 287.

The lesser Hop Trefoil, Decumbent Trefoil.

In Meadows — common. May to August. P.

TRIFOLIUM lupulinum minimum. R. S. 331.

TRIFOLIUM filiforme. H. F. 287.

Small Trefoil.

On gravelly Banks beyond South Street — uncommon.

May. A.

TRIFOLIUM pratense album. R. S. 327.

TRIFOLIUM repens. H. F. 283.

Dutch Clover, White flowered Meadow Trefoil.

In Fields and Meadows — very common. May to

September. P.

TRIFOLIUM pratense luteum capitulo Lupuli, vel agrarium. R. S. 330.

TRIFOLIUM agrarium. H. F. 286.

Hop Trefoil,

On Beacon Hill — not uncommon. June. A.

TRIFO-

TRIFOLIUM pratense purpureum. R. S. 328.

TRIFOLIUM pratense. H. F. 284.

Common Purple or Honey-suckle Trefoil.

In Fields and Meadows — very common. May to September. P.

TRIFOLIUM purpureum majus, folis longioribus
et angustioribus, floribus saturatioribus.
R. S. 328.

TRIFOLIUM medium. H. F. 284.

Long leaved Purple Trefoil with deeper coloured
Flowers.

*In Byfing Wood, and by Field sides in Ospringe —
not uncommon. July. P.*

TRIFOLIUM stellatum glabrum. R. S. 329.

TRIFOLIUM maritimum. H. F. 284.

Teasel-headed Trefoil.

On Shellness in Shepy — uncommon. June. P.

TURRITIS muralis minor. R. S. 294.

TURRITIS hirsuta. H. F. 254.

Hairy Tower Mustard.

On old Walls at Colkins in Boughton — very uncommon. May. B.

TUSSILAGO. R. S. 173.

TUSSILAGO Farfara. H. F. 315.

Common Colt's Foot.

In moist Fields — very common. April. P.

TYPHA.

TYPHA. RS. 436.

TYPHA *latifolia*. H. F. 345.

Great Cat's Tail, or Reed Mace.

On the sides of Ponds, and in Dykes — very common.
July. P.

TYPHA *palustris media*. R. S. 436.

TYPHA *angustifolia*. H. F. 345.

Narrow leaved Cat's Tail.

In the Ditches at Ewell and Nagden — not uncommon.
July. P.

V.

VALERIANA *sylvestris major*. R. S. 200.

VALERIANA *officinalis*. H. F. 12.

Great wild Valerian.

On the Banks of Ospringe River — not uncommon.
June. P.

VALERIANA *sylvestris minor*. R. S. 200.

VALERIANA *dioica*. H. F. 12.

Small wild or Marsh Valerian.

On the Banks of the shooting Meadow Stream —
common. June. P.

VALERIANELLA *arvensis præcox humilis*
semine compresso, sive Lactuca agnina.
R. S. 201.

VALERIANA *Locusta*. H. F. 12.

Lamb's Lettuce, or Corn Sallad.

In Corn fields, and rich cultivated Ground — common.
May. A.

VALE-

VALERIANELLÆ vulgaris species major, ferotina. R. S. 201.

VALERIANA *Locusta*. H. F. 13. β.

Late flowering Lambs Lettuce, or Corn Sallad.

In the Corn fields between Ore and Harty-ferry—not common. July. A.

VERBASCUM flore albo parvo. R. S. 287.

VERBASCUM *Lychnitis*. H. F. 76.

Hoary Mullein, or white Mullein.

On old Walls at Davington—not common. July. B.

VERBASCUM mas latifolium luteum. R. S. 287.

VERBASCUM *Thapsus*. H. F. 75.

Great white Mullein, High Taper, Cows Lungwort.

On dry Banks by Road sides—common. July. B.

VERBASCUM nigrum flore parvo, apicibus purpureis. R. S. 288.

VERBASCUM *nigrum*. H. F. 76.

Sage-leaved black Mullein.

On the sides of Chalk fields in Ospringe—very uncommon. As you ascend St. Martin's-Hill near Canterbury—plentifully. July. P.

VERBENA vulgaris. R. S. 236.

VERBENA *officinalis*. H. F. 505.

Vervain.

By Road sides and Foot paths—very common. July. P.

VERBE-

VERBESINA seu Cannabina aquatica flore minus
pulchro elatior. R. S. 187.

BIDENS tripartita. H. F. 309.

Trifid Water-hemp Agrimony.

In marshy watery Places — common. July. A.

VERBESINA pulchriore flore luteo. R. S. 187.

BIDENS cernua. H. F. 309.

Whole leaved Water-hemp Agrimony.

In Dykes near the Town Key — common. August. A.

VERONICA aquatica angustifolia minor. R. S.
280.

VERONICA scatellata. H. F. 5.

Narrow leaved Water Speedwell, or Brooklime.

In wet Ditches and marshy Ground — common. June. P.

VERONICA aquatica longifolia media. R. S. 280.

VERONICA Anagallis. H. F. 5.

The middle long leaved Water Speedwell, or
Brooklime.

In the Abby Ditches — common. July. P.

VERONICA aquatica rotundifolia Becabunga
dicta minor. R. S. 280.

VERONICA Becabunga. H. F. 4.

Common Brooklime.

*In wet Ditches and small running Waters — very
common. May. P.*

VERONICA

VERONICA Chamædrys sylvestris dicta. R. S. 281.

VERONICA Chamædrys. H. F. 5.

Wild Germander.

In Fields and Meadows — very common. May. P.

VERONICA floribus singularibus, in oblongis
pediculis, Chamædryfolia. R. S. 279.

VERONICA agrestis. H. F. 6.

Germander Speedwell, or Chickweed.

In Fields, on Rubbish, and in cultivated Ground —
common, May. A.

VERONICA flosculis singularibus Hederulæ folio,
Morsus gallinæ minor dicta. R. S. 280.

VERONICA hederifolia. H. F. 6.

Ivy leaved Speedwell, or small Henbit.

In dry Fields, and on old Walls — common. April. A.

VERONICA mas supina et vulgatissima. R. S.
281.

VERONICA officinalis. H. F. 4.

The Male Speedwell, or Fluellin.

In barren dry Fields — common. May. P.

VERONICA pretensis minor ; five Betonica Pauli.
R. S. 279.

VERONICA serpyllifolia. H. F. 4.

Little, or smooth Speedwell, or Paul's Betony.

In Fields and Meadows — common. May. P.

VIBURNUM. R. S. 460.

VIBURNUM *Lantana*. H. F. 112.

The pliant Mealy, or Way-faring Tree.

In Hedges — common. May. S.

VICIA. R. S. 320.

VICIA *saliva*. H. F. 278.

Common Vetch, or Tare.

In Fields among Corn — not uncommon. June. A.

VICIA *sepium perennis*. R. S. 320.

VICIA *sepium*. H. F. 278.

Bush Vetch.

In bushy Places and Hedges — common. May. P.

VICIA *minima præcox Parisiense*. R. S. 321.

ERVUM *soloniense*. H. F. 279.

Spring Tare.

In Ospringe Chalk Pits — common. April. P.

VICIA *sylvestris* five *Cracca major*. R. S. 321.

VICIA *latbyroides*. H. F. 279.

Strangle Tare, or Wild Vetch.

In sandy Fields in Hernhill — not uncommon.

May. A.

VINCA

VINCA *Pervinca minor*. R. S. 268.

VINCA *minor*. H. F. 77.

Periwinkle.

In Cades Wood, and Hedges in Preston and Ospringe
— *common* April. P. With a white Flower,
near the 39th Mile Stone — *very uncommon*.

VIOLA *bicolor arvensis*. R. S. 366.

VIOLA *tricolor*. H. F. 331. β .

Pansies, Heart's-ease, Two Faces under a Hood.
In sandy Corn fields at Hernhill — *common*. Au-
gust. A.

VIOLA *martia inodora sylvestris*. R. S. 364.

VIOLA *canina*. H. F. 331.

Wild or Dogs Violet.

In Woods and under Hedges — *very common*. April. P.

VIOLA *martia hirsuta inodora*. R. S. 365.

VIOLA *hirta*. H. F. 330.

Hairy Violet.

In the new Wood at Cades in Ospringe — *plentifully*.
March. P.

VIOLA *martia alba*. R. S. 364.

VIOLA *odorata*. H. F. 330. β .

White Violet.

Under dry Hedges and Banks, by Way sides — *not un-*
common. March. P.

VIOLA *maritima purpurea*. R. S. 364.

VIOLA *odorata*. H. F. 330.

Purple sweet Violet.

Under Hedges, and in Fields — very common.

March. P.

VIOLA *tricolor*. R. S. 361.

VIOLA *tricolor*. H. F. 331.

Pansies, Three Faces under a Hood.

In Corn fields at Ore, and in Gardens — common.

June. A.

VIRGA *aurea*. R. S. 176.

SOLIDAGO *Virga aurea*. H. F. 318.

Common Golden Rod.

In Woods and Hedges — common. August. P.

VISCUM. R. S. 464.

VISCUM *album*. H. F. 367.

Mistle, or Mistletoe.

On Trees, especially in Orchards — common.

May. S.

VULNERARIA *rustica*. R. S. 325.

ANTHYLLIS *Vulneraria*. H. F. 273.

Kidney Vetch, or Ladies Finger.

About Byfing Wood — common. July. P.

U.

ULMARIA. R. S. 259.

SPIRÆA *Ulmaria*. H. F. 191.

Meadow sweet.

In moist Meadows in Davington—common. July. P.

ULMUS folio glabro.

ULMUS *glabra*. H. F. 95. β.

The smooth leaved or Wych Elm.

In Hedges—not uncommon. April. S.

ULMUS folio latissimo scabro. R. S. 469.

ULMUS *glabra*. H. F. 95.

The Wych Hazel, or broad-leaved Elm.

In Hedges at Throwleigh—uncommon. April. S.

ULMUS vulgatissima, folio lato scabro. R. S.

468.

ULMUS *campestris*. H. F. 94.

The common Elm.

In Hedges—very common. April. S.

URTICA minor. R. S. 140.

URTICA *urens*. H. F. 355.

The lesser Stinging Nettle.

In cultivated and uncultivated Ground—very common.

August. A.

M 3

URTICA

URTICA racemifera major perennis. R. S. 139.

URTICA dioica. H. F. 355.

The common Stinging Nettle.

Under Hedges, and on Rubbish — very common
July. P.

PLANTS OMITTED.

* CALENDULA officinalis. Lin. Syst. Nat. II.

P. 577.

Garden Marigold.

On the Beach at St. Margaret's at Cliff; plentifully.

And in a Field at Woodnesborow — very
uncommon. June. A.

CRAMBE maritima brassicæ folio. R. S. 307.

CRAMBE maritima. H. F. 260.

Sea Colewort.

At St. Margaret's at Cliff — uncommon. June. P.

HYDROCERATOPHYLLON folio aspero quatuor cornibus armato. R. S. 135.

CERATOPHYLLUM demersum. H. F. 357.

Horned Pond-weed.

In Marsh Dykes — Common. June. P.

* MYAGRUM. R. S. 302.

MYAGRUM sativum. H. F. 242.

Gold of Pleasure.

Near Sandwich, among the Flax — plentifully.

June. A.

TRI-

TRIFOLIUM luteum lupulinum. R. S. 331.

MEDICAGO *lupulina*. H. F. 288.

Melilot Trefoil.

*In Paths near Westwood — not common. May to
August. A.*

CAUCULIS nodosa echinato semine. R. S. 220.

TORDILIUM *nodosum*. H. F. 98.

Knotted Parsley.

*By the Road side at Minster in Shepey — not common
May. A.*

LINARIA antirrhinum dicta. R. S. 283.

ANTIRRHINUM *minus*. H. F. 238.

The least Toad Flax.

*In a Corn field near Whitehill, Selling — uncom-
mon. A.*

SENECIO minor latiore folio, five montana.
R. S. 178.

SENECIO *sylvaticus*. H. F. 316.

Mountain Groundsel.

In Badgen Wood — not common. August. A.

OENANTHE aquatica triflora. R. S. 220.

OENANTHE *fistulosa*. H. F. 104. β .

The lesser Water Dropwort.

In Ditches at Graveney — common. July. P.

EMPHOTILUM...
MUSCICAGO...
...
...
...

CAS...
...
...
...

...
...
...
...

...
...
...
...

...
...
...
...

FOSSILIA SHEPEIANA.

A
SHORT VIEW
OF THE
FOSSIL BODIES
Native and Extraneous,
OF THE
ISLAND
OF
SHEPEY,
IN THE
COUNTY OF KENT.

Quicquid sub terra est in apricum proferet ætas.

INTRODUCTION.

THE Island of Shepey, with the annexed Peninsulas of Harty and Elmly, surrounded by the Waters of the Thames, the Medway, and the Sea, is upwards of Thirty Miles in Circumference, about Thirteen in Length from Shellness to Sheerness, and in its greatest Breadth about Six. It hath long been noted for producing large Quantities of Sheep (from whence probably it derived its Name) as well as Corn; and exhibits to the inquisitive Naturalist a most desirable Spot, by affording on its Surface many rare Plants, and more especially in the Bowels of its Northern Cliffs, so great a Quantity and Variety of Fossils, both native and extraneous, as are scarcely to be paralleled.

These Cliffs are in Length about Six Miles from East to West, gradually declining at each End; the more elevated Parts whereof reach about two-thirds of their extension, and are, at the very highest of them about Minster, not less than Thirty Yards in perpendicular Height above the Beach or Shore, and being composed of Clay, and constantly washed at their Basis by the
Tides

Tides which beat against them more especially when driven by strong Easterly Winds, they are continually wasting more or less, and falling down upon the Shore, where the Clay being by degrees washed away, the Fossils are left in great Abundance. So great is the Loss of the Land at the highest Parts, that sometimes near an Acre of it hath sunk down at once, though these Falls are indeed extraordinary.

Minster, Shurland, and Warden, are the three Manors to which these Cliffs appertain, the Proprietors whereof hire them out to different Owners of Coperas Works, who employ the neighbouring Poor to collect the Pyrites or Copperas Stone upon the Beach, at the Rate of about a Shilling per Bushel, which they deposit in Heaps, when collected, until a Quantity sufficient to load a Hoy is procured, when it is measured and paid for.

These Coperas Gatherers, from the Author's first Visits were induced to preserve and carry home every Thing that seemed to their Apprehension somewhat particular, and having since met with Encouragement, and acquired some Information relative to what was most desirable by the Curious, they have generally from that Time reserved such, until Inquiries have been made for them. Many rare Specimens, both native and extraneous, far beyond any equal Space of Ground, perhaps

perhaps the Universe can produce, are by these means preserved. Indeed, more may thus be obtained, at a very easy Expence, than by the most diligent Fossilists alone, as they cannot allow due Time to examine the Shore sufficiently. This is evidently confirmed by comparing the Specimens from this Coast, recited in the Works of our most celebrated * Naturalists, with those in the ensuing Catalogue. The Author doubts not that many curious Specimens have escaped his Inquiries, and as the Cliffs are daily dropping their Contents more or less, he is fully assured that there will always remain a Variety sufficient to gratify every Lover of the Study, whenever they may choose to take a Trip thither. But alas! One disagreeable Circumstance attending a considerable Part of the Fossils here collected, is, that they are so much impregnated with Pyritical Matter, that after being for some Time placed in a Cabinet, the Salts thereof shoot and entirely destroy them. Happy would it be, could some certain Remedy be discovered whereby this Accident might be prevented. The Loss of many valuable Specimens by this Cause, together with his Distance from any inquisitive and able Naturalists, at last induced the Author to dispose of his whole Collection to INGHAM FOSTER, Esq;

* Dr. Grew, Mr. Llhuyd, Mr. Petiver, Dr. Woodward, Mr. Dale, and Dr. Hill.

Merchant of London, where it now forms no inconsiderable Part of that Gentleman's very valuable Cabinet.

He, many Years since, entertained Thoughts of publishing a descriptive Catalogue of all the Fossils there, by him collected, but was prevented, as well by more necessary Attachments as the Difficulty of the Undertaking: However, it having been intimated to him that a concise Account of them would not be deemed an unacceptable Appendix to his PLANTÆ FAVERSHAMIENSES, in which several of the Shepey Plants are recited, as no special Account of these Fossils hath been offered to the Public, and as those that have been already noted by the before cited Authors are dispersed in their Works, and as many others have not as yet been published, he complied with the Proposal, and set about revising his Notes and Observations, and now offers this short Account of the Discoveries made there in his annual Visits for above these Thirty Years past; at first solely for the Purpose of collecting these elegant Bodies, and afterwards to superintend his private Affairs, having acquired some Property in the Island, not far distant from the Cliffs.

What the candid Reader may expect in the following Catalogue is, a general Account of the Variety of the Fossil Bodies, both native and extraneous, reduced into

some Order under particular Heads, with Notice of their degrees of Rarity. If what is offered shall afford any Information worthy the attention of the Lovers of Natural History, the Author will think himself sufficiently rewarded for this Attempt of exhibiting the Out-lines of a Work which merits the Abilities of a Person of more Judgment and greater Leisure, to enlarge and perfect, than he pretends to be possessed of. He cannot conclude however, without observing, that from the contemplation of so great a Variety of extraneous Fossils discovered in these Cliffs, which were evidently the produce of very different Climates, he thinks himself rationally induced to believe, that nothing short of an universal Deluge could be a Cause adequate to such an Effect.

E. J.

FOSSILIA SHEPEIANA.

NATIVE FOSSILS.

THE CLIFFS.

THESE are composed of three different Strata of brown Clay; the uppermost is of the lightest Colour, and most friable; the lowest darkest, and of a closer Texture.

PYRITÆ.

Copras Stones, are the most frequent of all the Fossils of the Cliffs: They are of various Forms and Sizes; round, botryoide, flat, varying in every Degree, as well in Figure as Colour, and impregnating most of the other Fossils found here.

LUDUS HELMONTII:

The Waxed Vein; Nodules of which are lodged in the Cliffs in great Numbers; they are generally flat and oval, and of various Sizes, sometimes exceeding two Feet in length, all are covered with Coats

in the Cliffs, but are frequently seen denudated on the Beach; the *Tali* are of different Sizes and Forms, generally they pass strait from one flat Surface of the Nodule to the other; in some there is a double Set of them, running only half through the Body, with intermediate sparry Partitions; which being generally of a waxen Colour, at first gave this Nodule its Name. The Number of the *Tali* in different Nodules are indeterminate; both the *Tali* and *Septa* vary in Colour in different Specimens — they are very common.

BEZOAR MINERALE.

Bodies, which are cruusted, ferrugineous, and of a rounded Form, covered with various Crusts of the Colour of the Clay of the Cliffs, and of different Sizes — These are common.

CALCULI.

Pebbles. Amongst the numbers on the Beach, there have been some Specimens collected, which by polishing have proved very elegant.

SELENITES.

Selenite, is found here in great Plenty, chiefly of the columnar sort, some Specimens exceed six Inches in length, but most are of shorter dimensions,

sions, which are common. The rhomboidal Sort is not near so common.

LEPASTRUM. D. HILL.

The Starry Waxed Vein, is a Fossil peculiar to these Cliffs, not having as yet been discovered elsewhere to the Author's Knowledge. Dr. Hill hath made from their Forms three Sorts of it, to which others might be added if nicely attended to. It is ever affixed to the Septa of the Ludus Helmontii, where the Fissures are so far divided as to permit the Stars to shoot. The largest Specimens do not exceed three Inches in Length, and these are very rare; those are generally of a much less Size that are common.

SUCCINUM.

Amber, is very uncommon; a few Specimens only, and those small and transparent, are to be found here.

EXTRANEOUS FOSSILS.

VEGETABLES.

LIGNUM FOSSILE.

Fossil Wood, in large Masses, seemingly of Oak from its Grain, even to above three Feet in Length,

is plentifully dispersed on the Beach, and generally impregnated with pyritical Matter.

EQUISETUM.

Horse-tail, Joints of the naked Species, are sometimes found even to the Length of three Inches, but they are uncommon.

FRUCTUS VARII.

Fossil Fruits, of very different Kinds, are here collected, in Variety and Number exceeding all that have been mentioned by Writers on this Branch of Natural History: Many of them have been drawn and compared with recent ones, in a particular Account given of them by the Author's late learned Friend, Dr. *James Parsons*, in the *Philosophical Transactions* — Vol. L. Part I. p. 396. These have by some been imagined mere Sports of Nature amongst the *Coperas*; but as all the Specimens have not the misfortune to be impregnated by it, some few, and those very elegant ones, being entirely free from it, and petrified, that Conceit without further Discussion, must fall to the Ground.

ARISTÆ.

Ears of Corn, some small Species, are sometimes to be met with, imperfect and impregnated with the *Pyrites*, but they are not common.

MYCE-

MYCETITÆ.

The Mushroom Stone. Here are two Kinds, the Conoide and Discoide, both small, and the first is by far the most frequently to be found.

ANIMALIUM PARTES.

ELEPHAS.

The Elephant. A considerable Number of the Bones of this very large Animal have been discovered on the Eastern Shore, a Mile from the Cliffs, on the Author's Manor of *Nutts*, in *Laysdown*; an Account of which may be seen in the Philosophical Transactions — Vol. *xlvi*. part *ii*. page 626.

TESTUDO.

The Tortoise. Two varieties have been found under the Cliffs, one of a more convexed Form, the other more depressed; both are extremely rare.

PISCIUM PARTES.

The Heads of Fish, from a very small Size to that of a middling Cod, are not unfrequent.

Palates of Fish, some oval, others of a squarish form, when perfect, or nearly so, are very uncommon;

common; but small Portions of them, especially of the square Sort are common: To these last, Mr. *Llbuyl* hath given the whimsical Name of *Scopula littoralis*, and to those of the first kind, *Bufonitæ*. In the Gentleman's Magazine for May, 1755, p. 409, is a Plate exhibiting both Kinds, in fine Preservation.

Teeth of Fish, usually called *Sharks Teeth*, are plentifully met with, except those of a large Size, which are very uncommon.

Back Bones of Fish, are very common in separate Joints, but with three or more united, are uncommon.

Parts of the Bodies of Fish, with the Scales on them, are not very uncommon, but none entire have come to the Author's Knowledge.

Tails of Fish, are frequently collected, and some few with a Vertebra united to them; there are very few of a large Size.

CANCRI MACROURI.

Lobsters, are not uncommon, but none of the Specimens are perfect; the Bodies and Tails united are not so frequent as when separated; the Claws are also found separately of various Sizes, but generally much injured.

Prawns

Prawns, are very rare; they are nearly perfect, and considerably larger than those taken recent on the opposite Coast of the Isle of *Thanet*.

CANCRI BRACHYURI.

Fossil Crabs. Before the Publication of Dr. *Hill's* History of *Fossils*, these were extremely rare in *England*; but since that Time, thousands of Specimens here collected, have passed through the Author's Hands. There are two or three Varieties, but all common except very fair Specimens, which are still very rare; their Size varies from that of a small Bean, to that of the common *Crab* of the Coast; some of them, especially the smaller ones, are impregnated by the *Pyrites*.

ECHINITÆ.

Echinites, here are the *Ovarius*, *Galeatus*, *Pileatus*, *Cordatus*, and *Placentiformis*; but all of them are very rare, especially the last.

The Spines, are of the foliated Kind, and that Sort named by Mr. *Elbuid*, *Volvola*—both are rare.

A S T E R I A S.

The Star Fish of five Rays, has not yet been found entire, and its single Rays are uncommon.

ASTROITÆ

ASTROITÆ ET ASTEROPODIUM.

Star Stones, with depressed Rays, and the Roots of them are very uncommon; the Specimens of the latter Kind were so imperfect, that without better Information the Author was at a Loss how to place them in his Collection.

BELEMNITES.

Belemnite, only one Specimen, and that imperfect, hath hitherto been discovered.

CONCHYLIA.

TUBULI MARINI.

Worm Shells, of various contorted Forms, generally of a small Size, some with their Shells, others denudated, frequently impregnated with the Coperas, are common.

LAPIS SYRINGOIDES.

The piped Waxen Vein, externally appears like the *Ludus Helmontii*, being found in incrusted Nudoles; they are evidently Oak that had first been eroded by Sea Worms, and afterwards their Perforations filled with Pyritical Matter and Spar,
similar

similar to the *Waxen Vein*, surrounding each Perforation ; these are very common.

CONCHYLIA UNIVALVIA.

DENTALIA.

The Tooth Shells, generally small, about an Inch long, curving towards the Apex in the more perfect Specimens, some covered with their Shells, and others denudated ; both Sorts are common.

PATELLA.

The Limpet, is a very rare Fossil, having only once been collected ; the Specimen was small, of the striated Sort, and somewhat imperfect.

NAUTILUS.

The chambered Nautilus, is a common Fossil of these Cliffs. These vary in Size from that of Hazel-Nut to that of a Man's Head, most commonly they have their pearly Shell, and are often immersed in Nodules ; there are also some small ones found of a ferrugineous Colour ; very few of the large ones are perfect, having received Injury by compression.

ORTHO-

ORTHOCERATITES.

Small Fragments seemingly of this *Genus* are found here, but very uncommon; the Chambers or Partitions appear to be united by a Dove-tail Suture.

TUBULI CONCAMERATI.

Cylindrical Bodies, with several Concamerations passing transversely through them, are generally of the Size of a Swan's Quill, and about two or three Inches in Length, some petrified, others wholly pyritical, and are not uncommon.

COCHLEÆ.

Snail Shells, with a short as well as with a produced Clavicle, some with their Shells, others denuded, are generally of a small Size, and impregnated by the *Pyrites*, and are uncommon.

TROCHI.

The Topshells, are of different Forms, some flatter, others more elevated, of various Sizes, some with Shells on, others denuded, most are impregnated with *Pyrites*, and are common.

CONCHA

CONCHA VENERIS.

The Cowry. This is small, denudated, pyritical, and extremely rare.

BUCCINA.

Whilkes, are found in considerable Quantities, varying in Form and Size, but generally small.

The Opercula are sometimes, though rarely seen.

NERITÆ.

The Nerites, are common; and generally impregnated with Pyrites.

TURBINES.

The Screw Shells, and casts of them without the Shell, of various Sizes, and mostly impregnated with Pyrites, are very common.

CONCHYLIA BIVALVIA.

OSTREA.

Fossil Oysters; generally of a small Size, are found here, but very uncommon; although the whole Coast is covered with recent ones.

O

CONCHÆ

CONCHÆ ANOMIÆ.

These Shells in great variety, tho' generally of a small Size, and impregnated with the Pyrites, are very common.

PECTUNCULI.

Cockles, smooth, striated, and fasciated, of a small Size, are found in tolerable Plenty; but fair and large Specimens are uncommon.

TELLINÆ.

The Wedge Shell. Small Specimens of which, generally denudated and pyritical, are very common.

MUSCULI.

Muscles. These are extremely rare, a few large Specimens, and one only of the common Muscle have been hitherto met with; although the Coast abounds with the latter.

PINNÆ.

The Sea Wing. Only small Fragments of this Shell have as yet been discovered, and these are very rare.

F I N I S.

I N D E X

OF THE

ENGLISH NAMES.

A.

Page

Abele Tree	87
Adder's Tongue	73
Agrimony	2
Dutch, or common Hemp	36
Alder, common	3
black Berry-bearing	39
Alexanders	108
Alkanet, evergreen	15
bastard	16
Ale-hoof	17
All-good	14
All-heal, Clowns	106
narrow leaved	106
All-feed, upright Blite	25
least Rupture-wort	92
Anemony Wood	6
Angelica, wild	6
Apple-Thorn	108
Archangel, red	53
yellow	54
white	52
Arsmart,	

	Page
Arsmart, common	82
dead or spotted	81
greater, with pale leaves	82
with spotted stalks	81
willow-leaved, perennial	82
Ash Tree	39
mountain	110
Ash-weed	6
Asp, or Aspen	87
Avens, common	21
water	22
B.	
Ba'm, garden	67
Basil, wild	2
great wild	28
Base Rocket	95
Batchelor's-buttons	91
Bear's-foot	45
Bedstraw, Ladies-yellow	40
Beech Tree	37
Beet, sea	12
Bell-flower, lesser round leaved	18
Berberry Bush	12
Betony, water	103
wood	112
Eindweed, black	37
great	28
small	28
sea	28
Birch Tree	13
Bird's Foot	77
trefoil	62
the greater	61

INDEX.

iii

Page

Bird's Nest, or Wild Carrot	32
Bitter-sweet	108
Blackberry	98
Blite, late flowering, or Fig leaved	13
the least	14
maple leaved	26
upright	26
round-leaved, or All-feed	25
sea	14
Blood wort	55
Borage	15
Bottle, blue	31
white	64
Box Tree	17
Brakes, common	38
Bramble, or Briar, common	98
small	98
Briar, sweet	97
Briony, black	112
white	15
Broom, common	40
rape	78
Brooklime, common	120
middle, long leaved	120
narrow leaved	120
Buckbean	67
Buckthorn	96
sea	96
Buck's-horn plantain	84
Buck-wheat, or Brank	37
Bugle	16
Bugloss, small wild	15
vipers	33
English vipers	33
Bullace	

	Page
Bullae Tree, black	90
white	90
Burdock	56
with smaller heads	56
taller, with a larger flower	56
with wooly heads	57
Burnet	101
Burnet-faxifrage, great	83
lesser round leaved	84
small	84
Bur-reed, great branched	110
not branched	110
Bush Vetch	122
Butcher's Broom	99
Butter-bur, common	82
Butter-cups	93

C.

Cabbage, sea	15
Calamint, common	18
Calf's-snout	7
Caltrops water, greater	88
lesser	88
Cammock	7
Camomile, sweet-scented creeping	25
Campion-bladder	64
corn, white	64
greater corn	64
red flowered	64
wild white	64
Dover	62
sea	62
narrow leaved	63
	Can-

I N D E X

v

Page

Canterbury Bells	18
Carrot, wild	32
Cat-mint	118
Car's-tail, great	118
lesser	118
Celandine, greater	79
lesser	25
Centory, lesser	24
lesser, with a white Flower	24
yellow	23
Charlock, white	96
or white Mustard	95
yellow	95
Cheese-rening	40
Chier, wild	59
Cherry wild, common	24
Cherry-wood	73
Chervil, small Hemlock	71
wild	24
Chesnut Tree	23
Chick-weed, breakstone	5
least	4
marsh, great	4
mouse-ear, broader leaved	3
narrow leaved	3
least	24
sea	4
purple flowered	111
star-headed	111
Chickweed or Speedwell	121
Cicely, wild	27
Cinque-foil, common	81
Tormentil	80
	Cistus,

	Page
Cistus, dwarf, or little Sun-flower	44
Clary, common English wild	47
Climber, great wild	27
Cleavers	7
Clot-bur	56
with smaller heads	56
taller, with a purple flower	56
with wooly heads	57
Clover, Dutch	116
Clown's All-heal	106
Cockle	63
Cock's comb, great marsh, red rattled	80
red rattle	80
yellow	80
Cock's-head	73
Codlings and Cream	65
Colewort, sea	126
Colt's-foot, common	117
Columbines	8
Comfrey	112
Corn, bottles	31
Feverfew	25
Marygold	26
Mint	68
Parsley, small	23
Sallad	118
late flowering	119
small	52
Cornel, female	29
Cotton Thistle	21
Cow's Lung-wort	119
Cow Parsnep	111
Cowslip, great	89
common	89
	Cow-

I N D E X.

vii

	Page
Cow-weed	27
Cow-wheat, common, wild	67
Crab Tree	66
Crane's-bill, dove's-foot	41
field	41
hemlock leaved	41
jagged leaved	42
long stalked	42
small flowered	42
sea	42
Cresses, water	107
winter, or Rocket	35
Crimson Grass Vetch	23
Cross-wort	30
Crowfoot, common creeping	94
corn	93
finest leaved	92
ivy leaved	93
round, rooted, or bulbous	93
sweet-wood	94
water fennel leaved	92
fine leaved	93
round leaved	94
various leaved	93
upright meadow	94
Crow-garlick	3
Cuckow flower	63
pint	8
Cudweed, black headed	43
common	43
least	43
long leaved, low branched	43
upright	43
	Cur-

	Page
Currants, black	96
red	97
D.	
Daffodil, wild English	71
Daifie, common	12
greater	58
Dandelion	32
rough, or Dandelion Hawkweed	32
Deptford Pink	22
Devil's-bit	102
yellow	47
guts	31
Dewberry	98
Dittander	58
Dock, common broad leaved	56
great water	55
greener leaved	56
red leaved	55
fiddle	55
sharp pointed	54
leffer	55
with curled leaves	55
Dodder	31
Dogberry	29
Dog's Mercury	69
Rose, red flowered	144
white flowered	143
Violet	123
Dove's-foot, common	41
jagged leaved	42
small flowered	42
Dropwort	37
Drop-	

I N D E X.

ix

	Page
Dropwort Hemlock	73
water	73
Duck's-meat	58
greater	58
ivy leaved	58
Dwale	12
Dwarf Elder	101
Dyer's Weed	41

E.

Earth-nut	16
Eglantine	97
Elder, common	100
Dwarf	101
water	73
Elecampane	44
Elm, broad leaved	125
common	125
smooth leaved, or Wych	125
Enchanter's Nightshade	27
Eryngo, Sea Holly	35
Eyebright	36
red	36

F.

Fennel, common	38
Hog, or Sulphur-wort	38
Fern, common male	38
female, or common brakes	38
small branched stone	38
Feverfew	67
corn	25
field	25
Figwort,	

x

I N D E X.

	Page
Figwort, common knobby rooted	103
water	103
Finkle	38
Five leaved grass	81
Flax, purging or wild dwarf	61
narrow leaved wild	60
perennial blue	61
Fleabane, middle	29
small	29
blue flowered	9
Flix-weed	36
Flower-de-luce, yellow water	51
Fluellin, round leaved	60
sharp pointed	60
Fool's Parsley	27
stones, female	75
male	75
Foxglove, purple	32
French Mercury, male and female	69
Frog-bit	112
lettuce	88
Fumitory	39
great white	39
Furze	41
needle	40

G.

Gentian, dwarf autumnal	41
Germander-speedwell, or Chickweed	121
wild	121
Gilli-flower, sea	111
Gladdon, stinking	51
	Glass.

I N D E X

xi

	Page
Glass-wort, white	14
jointed	51
prickly	51
Goat's-beard, purple	114
yellow	114
Gooseberry	44
Gold of Pleasure	126
Goldilocks	94
Golden rod, common	124
Goose-foot, common	14
sharp leaved	14
Goose-grass	7
marsh	7
Goose Tongue	91
Gorse	41
Goule	40
Grass-vetch, crimson	23
Green-wood	41
Go to bed at noon	114
Gout-weed	6
Gromwell, or Gromile	61
Ground Ivy	17
upright	106
Groundfel, common	105
mountain	127

H.

Hare bells	48
Hare's ear, the least	16
Hare's foot	115
Hart's tongue	83
Hawk nut, or Hog-nut	16
Hawk-	

	Page
Hawkweed, with bitten roots	47
bushy	46
narrow leaved bushy	46
long rooted	46
smooth succory	47
rough succory	47
Hawthorn	69
Hazel-nut Tree	29
Hazel-wych	125
Heart's-ease	123
Three faces under a hood	124
Heath common, or Ling	35
cross leaved	34
fine leaved	34
Pea, or wood pea	78
Hedge mustard	35
nettle	40
parsley	23
Hellebore, great bastard black	45
wild black	44
broad leaved bastard	45
white flowered bastard	45
Hell-weed	31
Hemlock	26
Hemlock Chervil, small with rough leaves	71
dropwort	73
lesser, or Fool's parley	27
water	83
Hemp-agrimony, common or Dutch	36
Water trifid	120
whole leaved	120
Henbane, common	48
Henbit, great	53
small	121
Hep-	

I N D E X.

xlii

Page

Hep-tree	97
Herb, Bennet, or common Avens	21
Gerard	6
Paris	45
Robert	42
Two-pence	72
High-taper	119
Hog-fennel	83
Hog-nut	16
Holly-tree	2
Hone-wort	107
Honeyfuckle, common	18
trefoil	117
Hop, male and female	62
trefoil	116
Horehound, stinking	11
water	64
white	66
Horn-beam Tree, or Horse-beech	78
Horse-radish	96
Horse-tail, corn	33
lesser, marsh	34
river	34
wood	34
Horse-shoe Vetch	37
Hound's-tongue, great	31
lesser green leaved	31
House-leek	104
Hyacinth, English	48
I.	
Jack-by-the-hedge	46
Iron-wort	106
b	Ivy,

	Page
Ivy, climbing or berried	44
Juniper, common	51
K.	
Kidney-vetch	124
wort	29
Knapweed, common	50
great	50
Knawel, annual	52
Knee-holly	99
Knot-grass, common	86
German	52
oblong narrow leaved	86
small narrow leaved	86
L.	
Ladies Bedstraw, mountain	70
white	70
yellow	40
Smock	18
small flowered	19
Finger	124
Thistle	20
unspotted	20
Traces, triple	77
Lamb's lettuce	118
late flowering	119
Lang-de-beuf	46
Larkspur, wild	32
Lavender, sea	60
Laurel, dwarf	57
Lettuce,	

I N D E X.

xv

	Page
Lettruce, Ivy leaved wild	52
least wild	52
Lilly-convally	59
water white	72
yellow	72
Ling	35
Live-long	5
Lords and Ladies	8
Loose-strife, greater smooth-leaved	64
least smooth-leaved coddled	65
lesser hairy coddled with small flowers	65
middle smooth-leaved	65
purple spiked	93
yellow	65
Loufewort, common	80
marsh, red rattle, Cock's-comb	80
Liquorice, wild or Vetch	9

M.

Madder, little field	98
great bastard	70
small mountain	70
wild, or great bastard	70
Madwort, German	19
Maiden hair, white	99
common black	2
English	115
Mallow, common	66
small wild, or dwarf	66
marsh	5
Maple, common	1
greater, or Sycamore	1

	Page
Mare's-tail	59
Marigold, corn	26
marsh	86
garden	126
Marjoram, pot	77
wild	77
Matfellow, common	50
great	50
May Lilly	59
weed, stinking	24
Meadow Pink	63
Rue	113
Sweet	125
Medick-Vetchling	73
Melilot	67
trefoil	127
Mercury, common English	14
dog's	69
French, male and female	69
Milfoil	69
water, feathered or spiked	87
verticillated	81
Milk-wort	85
sea	43
Mill-mountain	61
Mint, Nep or Nap	72
long leaved horse	68
round leaved horse	68
spear	68
water	68
with whorled coronets	68
Missel or Misseltoe	124
Money-wort	72
purple flowered	72
Moscatel,	

I N D E X.

xvii

Page

Moscatel, tuberous	70
Mother-wort	19
Mother of thyme, common	15
Mountain Ash	110
Mouse-ear scorpion grass	70
common creeping	83
water scorpion grass	71
Mug-weed	30
Mug-wort	8
Mullein, great white	119
hoary or white	119
fage leaved black	119
Musk Thistle	20
Mustard, common	106
hedge	35
tower, lesser	117
white	106
Myrtle, Dutch	40
N.	
Navel wort	29
Navew, wild	71
Needle Furze	40
Nep or Nap, or Cat-Mint	72
Nettle, dead	52
hedge	40
hemp-leaved, with a parti-coloured	
flower	53
white flowered	53
red	53
yellow	54
stinging, common	125
lesser	125
b 3	Nettle-

	Page
Nettle-hemp, white flowered	53
fair flowered	53
yellow	54
Nightshade, common	109
deadly	12
Enchanter's	27
woody	108
Nipple-wort	54
Nose-bleed	69
Nut-earth, Kipper, Pig, Hawk or Hog	16

O.

Oak Tree, common	91
with acorns on short-foot stalks	91
Oculus Christi, or Christ's eye	47
One Berry	45
Orach, common wild	13
grass leaved	10
narrow leaved wild	10, 11
with a spear pointed leaf	10
jagged sea	10
wild sea narrow leaved	10
wild sea	11
stinking	13
Orchis, female	75
female handed	76
male, or male fool's stones	75
male handed	76
butterfly	74
bee	74
bee hairy, or humble bee	77
fly	75
frog	76
	Orchis,

I N D E X

xix

Page

Orchis, man green	74
purple	75
Neapolitan	74
little purple flowered	76
purple late flowering	76
yellow or musk	75
bird's nest	74
ladies traces	77

Orpine	5
--------	---

Oslor	100
-------	-----

Ox-eye	88
--------	----

lips	89
------	----

tongue	46
--------	----

P.

Pagils, common	89
----------------	----

Pansies, two faces under a hood	123
---------------------------------	-----

three faces under a hood	124
--------------------------	-----

Park-leaves	49
-------------	----

hairy	48
-------	----

Parsley, garden	82
-----------------	----

corn or Honewort	101
------------------	-----

corn small	82
------------	----

hedge	82
-------	----

knotted	127
---------	-----

stone bastard	107
---------------	-----

Parsnep, Cow	111
--------------	-----

great water	107
-------------	-----

least water	101
-------------	-----

water creeping	101
----------------	-----

wild	79
------	----

Paul's Betony	12
---------------	----

Pea,	12
------	----

INDEX

Page		Page
57	Pea, everlasting broad leaved	57
57	narrow leaved	57
78	wood or heath	78
91	Pear, wild	91
79	Pellitory of the wall	79
91	bastard	91
91	Penny royal	91
48	Penny-wort, marsh	48
29	wall	29
58	Pepper-wort	58
82	Pepper, water	82
27	Periwinkle, the greater	27
23	the lesser	23
82	Pestilent Wort	82
40	Petty Whin	40
16	Pig nut or Hog-nut	16
25	Pilewort	25
5	Pimpernel, male	5
5	yellow	5
101	round leaved water	101
5	Pin-Pearl-wort	5
22	Pink, Deptford	22
63	meadow	63
12	Pipperidge Bush	12
84	Plantain, Buck's-horn	84
84	great	84
85	hoary	85
85	ribwort	85
85	sea	85
85	narrow leaved sea	85
84	water great	84
31	star-headed	31
86	Polypody	86
		Pond-

I N D E X.

xxi

Page

Pond-weed, broad leaved	80
fine or fennel leaved	88
grass leaved	88
horned	126
perfoliated	89
sea	88
Poplar Tree, black	87
white	87
trembling	87
Poppy, Corn	79
red	79
spatling	64
yellow horned	78
Plowman's spikenard	11
Prick-madam	104
Prick-wood, Cornel-tree	29
Spindle-tree	36
Primrose, common	9
Privet or Prim	59
Pudding-grass	9
Purslane, water	7
sea	11
Pursley Piert	18

Q.

Quick	69
Quicken Tree	110

R.

Radish, Horse	96
Water	92
Ragwort,	96

INDEX.

	Page
Ragwort, broad leaved marsh	50
common	50
hoary perennial with Groundfel	50
leaves	50
Ramsons	3
Raspberry	98
Rattle, yellow	80
red marsh	80
Reed Mace	118
lesser	118
Rest-harrow, Corn	7
spinous	7
creeping	6
Ribwort Plantain	85
Roan Tree	110
Rocket, or Winter Cresses	35
base	95
water	35
sea	7
Rose-bay, Willow herb	66
Rose, Burnet	97
dog's red flowered	97
white flowered	97
Rue, meadow	113
wall	99
whitlow grass	101
Rupture-wort, the least	96
Ruth, flowering	7
 Sage wood	103
Saint-foin	73
Saint	73

I N D E X.

xxiii
Page

Saint John's-wort	48
hairy	48
imperforate	49
least trailing	49
upright	49
Saint Peter's-wort	49
Sallow, common round leaved	100
Salt-wort, annual	99
black	43
Sampire	30
golden	9
marsh	99
perennial	81
Sanicle	101
Satyrium Royal, male handed	26
female handed	26
Sauce alone	26
Saw-wort	105
Saxifrage, common golden	101
meadow	105
white	102
Rue-leaved	102
Scabious, common field	102
lesser	102
Sheep's hairy	95
Scad Tree, or Scad Plumb	90
Scorpion-grass	20
water	71
Sea Bindweed	28
Buckthorn or fallow	96
Cabbage	15
Campion	62
Colewort	26
Pond-weed	88
	Sea

	Page
Sea Holly	35
Heath smooth	63
Gilly-flower	111
Rocket	17
Lavender	60
Plantain	85
narrow leaved	85
Spurrey, small flowered	110
Starwort	9
Scurvy-grass	28
Self-heal	90
Sengreen, white	103
common	104
Septfoil	114
Service Tree, common	69
Setter-wort	45
Shepherd's Needle	103
Purse	17
Rod	33
Silver Weed	80
Simson	105
Sloe Tree	90
Smallage	8
Snap-dragon, greater	7
Sneeze-wort	91
Snow-drop	59
Soap-wort, common	63
Sorb Tree	69
Sorrel, common	54
Sheep's	54
Wood	78
Sow-bane	14
Sow-	

I N D E X.

xxxv

	Page
Sow-thistle, common	109
prickly with jagged leaves	109
with less jagged leaves	109
smooth with less jagged leaves	109
Tree	110
Spear-thistle	19
Spear-wort, great	94
lesser	94
Speedwell, or Chickweed	121
Germander	121
ivy-leaved	121
little smooth	121
male	121
water, narrow leaved	120
with long leaves	120
Spikenard, Plowman's	11
Spatling Poppy	64
Spindle Tree	36
Spleenwort, or Miltwaste	9
rough	61
Spurge, broad leaved	114
dwarf, small annual	113
Laurel	57
petty	113
Sun, or Wart-wort	113
Wood	113
Spurrey, Corn	4
purple	111
sea	4
small flowered	110
Squinancy-wort	98
Star-headed Water Chickweed	111
Star of the Earth	84
Thistle	21
Star-	

	Page
Star-wort, autumnal	111
vernal	111
sea	9
Stich-wort, greater	22
least	5
lesser	22
middle, a variety	22
Stonecrop, yellow	104
Wall Pepper	104
yellow reflexed	104
mountain	103
insipid	104
Stone Parsley, bastard	107
Strangle Tare	122
Strawberry	38
barren	39
trefoil	115
Succory, wild	26
Sulphur-wort	83
Sun flower, little	44
Sweet Briar	97
Swine's Cress	71
Sycamore	1

T.

Tansey, common	112
wild	80
Tare, common	122
everlasting	57
hairy	30
small wild	30
smooth	30
spring	122
Tare,	

I N D E X.

xxvii

Page

Tare, strangle	122
tine	30
with smooth pods	30
Teasel, wild	33
small wild	33
Tent-wort	99
Thistle, common way, or creeping	21
Carline, dwarf	19
wild	21
Cotton	21
Marsh	20
Milk, or Lady's	20
without spots	20
Musk	20
Spear	19
Star	21
upon Thistle	19
Wetted, with small flowers	20
Thorn Apple	108
Thorn, black	90
white	69
common purging	96
sea buck	96
Thorough wax	16
Three faces under a hood	124
Thrift	111
Throat-wort, great	18
Thyme, mother of	105
Lemon	105
Tine Tare	30
with smooth pods	30
Toad Flax, common yellow	60
the least	127
Tooth-wort	6

	Page
Tormentil	114
Cinquefoil	80
creeping	114
Tower Mustard, hairy	117
Traveller's Joy	27
Tree, pliant mealy	122
Trefoil, Bird's foot	62
greater	61
common purple	117
Hare's-foot	115
Heart	115
Hedge-hog	115
Hop	116
lesser	116
long leaved purple	117
Marsh	67
Melilot	127
oval headed	116
Teasel headed	117
small	116
Strawberry	115
white flowered meadow	116
Triple Ladies Traces	77
Tutsan	49
hairy	48
Turnep	96
Tway-blade, common	13
Two faces under a hood	123
V.	
Valerian, great wild	118
jagged leaved	52
small wild or marsh	118
Venus-	

I N D E X.

xxix

Page

Venus-comb	103
Vervain	119
mallow	2
Vetch Bush	122
Crimson grass	23
common	122
tufted	30
wild	122
spring	122
tufted Horse-shoe	37
Kidney, or Lady's finger	124
Vetchling, common yellow bastard	57
Medick	73
Violet, Dog's or wild	123
purple sweet	124
Throat wort leaved	123
white	123
water	47
hairy	123
Viper's Bugloss	33
English	33
W.	
Wake Robin	8
Wall-flower	59
Penny-wort	29
Pepper	104
Rue	99
wort	101
Wart-wort	113
Water Betony	103
Caltrops, greater	88
lesser	88
Water	

	Page
Water Crow-foot, fennel leaved	92
fine leaved	93
finest leaved	92
Ivy leaved	93
round leaved	94
various leaved	93
Cresses	107
Dock, great	79
Dropwort	73
Hemlock	73
Elder	73
Figwort	103
Gladiole	17
Hemlock	83
Hemp agrimony, trifid	120
with an undivided leaf	120
Horehound	64
Lilly, white	72
yellow	73
Mill-foil, feathered or spiked	87
verticillated	81
Mint	107
hairy	107
with whorled coronets	88
Parsnep, creeping	108
great	107
least	101
Pepper	82
Plantain, great	84
star-headed	108
Purslane	77
Ragwort	50
Water	

xxxi

	Page
Water Speedwell, middle long leaved	120
narrow leaved	120
Wort	111
Violet	47
Way-bread	84
Way-faring Tree	122
Weed, yellow	62
Whin	41
petty	7
White-glaswort	14
White-rot	48
White-thorn	69
Whitlow-grafs, common	79
rue leaved	182
Williams, wild	63
Willow, common white	99
round eared shining	100
fand	100
sweet	40
Willow-herb, purple spiked	99
great flowered	65
greater smooth leaved	64
least smooth leaved coddled	65
lesser hairy coddled with small	
flowers	65
middle smooth leaved	65
Rose bay	66
yellow	65
Winter Cresses	35
Woad, wild	62
Wood · Anemone	6
Betony	12
bine	18
roof	8
	Wood

	Page
Wood Pea	78
Sorrel	78
waxen	59
Sage	103
Spurge	113
Wormwood, common	1
white sea	1
Wort, white glass	14
Wych Elm	125
Hazel	125
Y.	
Yarrow, common	69
Yellow-weed, common	62
Yew Tree	112
Willow, common	112
Willow, hard, purple jointed	112
Willow, great flowered	112
Willow, greater smooth leaved	112
Willow, least smooth leaved	112
Willow, lesser hairy coddled with small flowers	112
Willow, middle smooth leaved	112
Willow, Kotsch	112
Willow, yellow	112
Winter Cress	112
Wood, wild	112
Wood, Aromatic	112
Wood, decay	112
Wood, pine	112
Wood, root	112

I N D E X

O F T H E

L I N N E A N N A M E S

	Page
Acer campestre	1
Pseudo platenus	69
Achillea Millefolium	91
Ptarmica	70
Adoxa Moschatellina	6
Aegopodium Podagraria	27
Aethusa Cynapium	2
Agrimonia Eupatoria	63
Agrostemma Githago	16
Ajuga reptans	31
Alisma Damasonium	84
Plantago	3
Allium vineale	3
Ursinum	5
Althæa officinalis	14
Amarantus Blitum	5
Anagallis arvensis	15
Anchusa sempervirens	6
Anemone nemorosa	Anethum

	Page
Anethum Foeniculum	38
Angelica sylvestris	6
Anthemis Cotula	24
nobilis	25
Anthyllis Vulneraria	124
Antirrhinum minus	127
Elatine	60
Linaria	60
majus	7
spurium	60
Aphanes arvensis	81
Apium graveolens	8
Petroselinum	82
Aquilegia vulgaris	8
Arctium Lappa β . γ . δ .	56, 57
Arenaria marina β .	4
marina γ .	110
rubra	111
peploides	4
serpyllifolia	4
Artemisia Absinthium	1
maritima	1
vulgaris	8
Arum maculatum	8
Asperula odorata	8
Cynanchica	98
Asplenium Adiantum nigrum	2
Ceterach	9
Ruta muraria	99
Scolopendrium	83
Trichomanes	115
Astragalus glycyphyllos	9
Aster Tripolium	9
Atriplex	

I N D E X.

xxxv

Page

Atriplex hastata	10
hastata β .	11
laciniata	10
litoralis β .	11
litoralis	10
patula	10
ferrata	10
portula coides	11
Atropa Belladonna	12

B.

Ballota nigra	11
Bellis perennis	12
Berberis vulgaris	12
Beta vulgaris	12
Betonica officinalis	12
Betula alba	13
Alnus	3
Bidens tripartita	120
cernua	120
Borago officinalis	15
Brassica Napus	71
Rapa	95
oleracea	15
Blackstonia perfoliata	23
Bryonia alba	15
Bunium Bulbocastanum	16
Bunias Cakile	17
Bupleurum rotundifolium	16
tenuissimum	16
Butomus umbellatus	17
Buxus sempervirens	17

Calen-

Page		Page
10		
11		
10		
	C.	
	Calendula officinalis	126
	Callitriche autumnalis	111
10	verna	111
	Caltha palustris	86
	Campanula rotundifolia	18
11	Trachelium	18
	Cardamine parviflora	19
	pratensis	18
	Carduus acanthoides	20
11	acaulos	19
11	crispus	19
11	lanceolatis	19
11	Marianus	20
11	Marianus β.	20
11	nutans	20
3	palustris	20
	Carlina vulgaris	21
	Carpinus Betulus	78
	Caucalis Anthriscus	23
11	arvensis	23
	Centaurea Calcitrapa	21
11	Cyanus	31
23	Jacea	50
11	Scabiosa	50
	Cerastium aquaticum	4
11	semidecandrium	24
20	viscosum	3
10	vulgarum	3
	Ceratophyllum demersum	126
	Chierophyllum sylvestre	27
	temulum	24
		Cheiran-

I N D E X.

xxxvii

Page

Cheiranthus Cheiri	59
Chelidonium majus	79
Glaucium	78
Chenopodium album	13
Bonus Henricus	14
hybridum	26
maritimum	14
murale	14
polyspermum	25
rubrum	14
serotinum	13
Vulvaria	13
urbicum	26
Chrysanthemum Leucanthemum	58
segetum	26
Chrysofenium oppositifolium	101
Cichoreum Intybus	26
Circœa Lutetiana	27
Cistus Helianthemum	44
Clematis Vitalba	27
Clinopodium vulgare	28
Cochlearia anglica	28
Armoracia	96
Coronopus	71
Conium maculatum	26
Convallaria majalis	59
Convolvulus arvensis	28
sepium	28
Soldanella	28
Conyza squarrosa	11
Cornus sanguinea	29
Corylus Avellana	29
Cotyledon Umbilicus Veneris	29
Crambe maritima	126

d

Cratægus

	Page
Cratægus Oxyacantha	69
torminalis	69
Crepis tectorum	47
biennis	47
Crithmum maritimum	30
Cucubalus Behen	64
viscosus	62
Cuscuta europæa	31
Cynoglossum officinale	31
officinale β .	31

D.

Daphne Laureola	57
Datura Stramonium	108
Daucus Carota	32
Delphinium Consolida	32
Dianthus Armeria	22
Digitalis purpurea	32
Dipsacus pilosus	33
sylvestris	33
Draba verna	79

E.

Echium anglicum	33
vulgare	33
Epilobium angustifolium	66
hirsutum	65
montanum	64
palustre	65
ramosum	65
tetragonum	65
Equi-	

INDEX.

xxxix

Page

Equisetum arvense	33
fluviatile	34
palustre	34
sylvaticum	34
Erica cinerea	34
Tetralix	34
vulgaris	35
Erigeron acre	9
Ervum hirsutum	30
solonienſe	122
tetraspermum	30
Eryngium maritimum	35
Eryſinum Alliaria	46
Barbarea	35
officinalis	35
Euonymus europæus	36
Eupatorium cannabinum	36
Euphorbia Amygdaloides	113
exigua	113
Helioscopia	113
Peplus	113
platyphyllos	114
Euphrasia officinalis	36
Odontites	36

F.

Fagus Castanea	23
ſylvaticus	37
Ficaria verna	25
Filago germanica	43
montana	43
Fragraria ſterilis	39
veſca	38

	Page
Frankenia lævis	63
Fraxinus excelsior	39
Fumaria officinalis	39
claviculata	39

G.

Galanthus nivalis	59
Galeopsis Ladanum	105
Tetrahit	53
β.	53
δ.	53
Galeobdolon	54
Galium Aparine	7
uliginosum	7
verum	40
Mollugo	70
montanum	70
palustre	70
Genista anglica	40
tinctoria	41
Gentiana Amarella	41
Centaurium	24
Geranium cicutarium	41
columbinum	42
dissectum	42
molle	41
pusillum	42
maritimum	42
robertianum	42
Geum urbanum	21
rivale	22
Glaux maritima	43
Glechoma	

IX N D E X

xli

Page

Glechoma arvensis	106
hederacea	17
Gnaphalium sylvaticum	43
uliginosum	43
H.	
Hedera Helix	44
Hedysarum Onobrychis	73
Helleborus foetidus	45
viridis	44
Heracleum Spondylium	111
Hieracium Pilosella	83
fabaudum	46
umbellatum	46
Hippocrepis comosa	37
Hippophae Rhamnoides	96
Hippuris vulgaris	59
Hottonia palustris	47
Humulus Lupulus	62
Hyacinthus non scriptus	48
Hydrocharis Morfus ranæ	112
Hydrocotyle vulgaris	48
Hyoscyamus niger	48
Hypericum Androsæmum	49
hirsutum	48
humifusum	49
montanum	49
perforatum	48
pulchrum	49
quadrangulum	49
Hypochæris radicata	46

Ilex

Page

100

17

Jasione montana

Jlex Aquifolium

Inula crithmoides

dysenterica

Helenium

Pulicaria

Juniperus communis

Iris foetidissima

Pseudacorus

111

112

113

114

Lactuca saligna

Lamium album

amplexicaule

rubrum

Lapsana communis

Lathræa squamaria

Lathyrus latifolius

Nissolia

pratensis

sylvestris

Lemna minor

polyrrhiza

trifulca

Leontodon autumnale

hispidum

Taraxacum

Leonurus Cardiaca

Lepidium latifolium

Ligustrum vulgare

115

I.

H.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

Page

95

2

9

29

44

29

51

51

51

51

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

L.

Linum

	Page
Linum catharticum	61
perenne	61
Radiola	62
tenuifolium	60
Lithospermum officinale	61
arvense	16
Lithrum Salicaria	99
Lonicera Periclymenum	18
Lotus corniculata	62
γ.	61
Lychnis dioica	64
β.	64
Flos cuculi	63
Lycopsis arvensis	15
Lycopus europæus	64
Lyfimachia nemorum	15
tenella	72
vulgaris	65
Nummularia	72

M.

Malva Alcea	2
rotundifolia	66
sylvestris	66
Marrubium vulgare	66
Matricaria Chamomilla	25
inodora	25
Parthenium	67
Medicago arabica	115
arabica γ.	115
lupulina	127
Melampyrum sylvaticum	67
Melissa	

	Page
Melissa Calamintha	68
officinalis	67
Mentha aquatica	68
γ .	107
arvensis	68
longifolia	68
rotundifolia	68
spicata	68
Pulegium	69
Menyanthes trifolia	67
Mercurialis annua	69
perennis	69
Myrica Gale	40
Myosotis scorpioides	70
δ .	71
Myriophyllum spicatum	87
verticillatum	81
Myagrum fativum	126

N.

Narcissus Pseudo-Narcissus	71
Nepeta Cataria	72
Nymphæa alba	72
lutea	72

O.

Oenanthe crocata	73
fistulosa	73
β .	127
Ononis arvensis	7
repens	6
spinosa	7
Onopor-	

I N D E X.

xiv

	Page
Onopordum Acanthium	21
Ophioglossum vulgatum	73
Ophrys Anthropophora	74
apifera	74
apifera β .	77
Monorchis	75
Nidus avis	74
muscifera	75
ovata	13
spiralis	77
Orchis bifolia	74
latifolia	76
mascula	75
maculata	76
militaris	74
Morio	75
purpurea	75
pyramidalis	76
ustulata	76
Origanum onites	17
vulgare	77
Ornithopus perpusillus	77
Orobanche major	78
Orobis tuberosus	78
Osmunda spicant	61
Oxalis Acetofella	78
P.	
Papaver Rhœas	79
Parietaria officinalis	79
Paris quadrifolia	45
Pastinaca fativa	79
	Pedicu-

	Page
Pedicularis palustris	80
sylvatica	80
Peplis Portulaca	87
Pteris aquilina	38
aquilina β .	38
Peucedanum officinale	83
Phellandrium aquaticum	83
Picris Echioides	46
Pimpinella major	83
saxifraga	84
β .	84
Plantago Coronopus	84
lanceolata	85
Leoflingii	85
major	84
maritima	85
media	85
Polygala vulgaris	85
Polygonum amphibium	82
aviculare	86
β . et γ .	86
Convolvulus	37
Fagopyrum	37
Hydropiper	82
pensylvanicum	82
Perficaria	81
β .	81
Polypodium Filix mas	83
vulgare	86
Populus alba	87
nigra	87
tremula	87
	Poten-

I N D E X.

xlvii

Page

Potentilla Argentina	80
argentea	80
reptans	81
Potamogeton crispum	88
natans	89
marinum	88
gramineum	88
pectinatum	88
perfoliatum	89
ferratum	88
Poterium Sanguisorba	101
Prenanthes muralis	52
Primula veris α .	89
β .	89
vulgaris	89
Prunella vulgaris	90
Prunus Avium	24
insititia	90
insititia β . et γ .	90
spinosa	90
Pyrus Malus	66
communis	91

Q.

Quercus Robur	91
β .	91

R.

Ranunculus acris	94
aquatilis	93
aquatilis γ . δ .	92
aquatilis β .	93
arvensis	93

Ranun-

	Page
Ranunculus Auricomus	94
bulbosus	93
Flammula	94
hederaceus	93
Lingua	94
repens	94
sceleratus	94
Raphanus Raphanistrum	96
	95
Reseda lutea	95
Luteola	62
Rhamnus catharticus	96
Frangula	39
Rhinanthus Crista galli	80
Rosa arvensis	97
canina	97
Eglanteria	97
spinosissima	97
Ribes rubrum	97
nigrum	96
Uva crispa	44
Rubus cæsius	98
fructicosus	98
Idæus	98
Rumex Acetosa	54
Acetofella	54
acutus	54
Britannicus	55
crispus	55
obtusifolius	56
pulcher	55
sanguineus	55
Rumicis acuti varietas	55, 56
Ruscus	

I N D E X.

xlix

Page

Ruscus aculeatus

99

S.

Sagina erecta

5

procumbens

5

Salix alba

99

arenaria

100

amygdalina

100

capræa

100

vimenalis

100

Salicornia europæa β .

51

europæa

99

Salsola Kali

51

Salvia Verbenaca

47

Sambucus Ebulus

101

nigra

100

Samolus Valerandi

101

Sanicula europæa

101

Saponaria officinalis

63

Satyrion viride

76

Saxifraga trydactylites

102

granulata

102

Scabiosa arvensis

102

Columbaria

102

Succisa

102

Scandix Anthriscus

71

Pecten Veneris

103

Sedum acre

104

annuum

103

reflexum

104

reflexum β .

104

Telephium

5

sexangulare

104

e

Scleran-

	Page
Scleranthus annuus	52
Sempervivum tectorum	104
Senecio aquaticus	50
crucifolius	50
Jacobæa	50
sylvaticus	127
vulgaris	105
Serapias latifolia β .	45
latifolia	45
longifolia	45
Scrophularia aquatica	103
nodosa	103
Serratula arvensis	21
tinctoria	105
Seseli Caruifolia	105
Sherardia arvensis	98
Silene amœna	62
conoidea	63
Sinapis alba	106
arvensis	95
nigra	106
Sison Amomum	107
inundatum	108
segetum	107
Sisymbrium amphibium α .	92
Nasturtium	107
Sophia	36
sylvestre	35
Sium latifolium	107
nodiflorum	108
Smyrnum Olusatrum	108
Solanum Dulcamara	108
nigrum	109
Solidago Virga aurea	124
Sonchus	

I N D E X.

li

Page

<i>Senecio arvensis</i>	110
<i>oleraceus</i> <i>α. β. γ. δ.</i>	109
<i>Sorbus aucuparia</i>	110
<i>Sparganium erectum</i> <i>β.</i>	110
<i>Spartium scoparium</i>	40
<i>Spergula arvensis</i>	4
<i>Spiraea Filipendula</i>	37
<i>Ulmaria</i>	125
<i>Stachys palustris</i>	106
<i>sylvatica</i>	40
<i>Stellaria graminea</i>	22
<i>graminea</i> <i>β.</i>	22
<i>Holostea</i>	22
<i>Statice Armeria</i>	111
<i>Limonium</i>	60
<i>Symphytum officinale</i>	112

T.

<i>Tamus communis</i>	112
<i>Tanacetum vulgare</i>	112
<i>Taxus baccata</i>	112
<i>Teucrium Scorodonia</i>	103
<i>Thalictrum flavum</i>	113
<i>Thlaspi Bursa Pastoris</i>	17
<i>Thymus Acinos</i>	2
<i>Serpyllum</i>	105
<i>Serpyllum</i> <i>α.</i>	105
<i>Tordylium nodosum</i>	127
<i>Tormentilla erecta</i>	114
<i>reptans</i>	114
<i>Tragopogon pratense</i>	114
<i>porrifolium</i>	114
c 2	
<i>Trifolium</i>	

	Page
Trifolium agrarium	116
arvense	115
filiforme	116
fragiferum	115
medium	117
Melilotus officinalis	67
maritimum	117
pratense	117
procumbens	116
repens	116
scabrum	116
Tussilago Farfara	117
Petasites	82
Turritis hirsuta	117
Typha latifolia	118
angustifolia	118

V.

Valantia Cruciata	30
Valeriana dioica	118
officinalis	118
Locusta	118
Locusta α .	52
Verbascum Lychnitis	119
nigrum	119
Thapsus	119
Verbena officinalis	119
Veronica agrestis	121
Anagallis	120
arvensis	121
Becabunga	120
Chamædrys	121
hederifolia	121

Veronica

I N D E X.

lii

	Page
Veronica officinalis	121
serpyllifolia	121
scutellata	120
Viburnum Lantana	122
Opulus	73
Vicia Cracca	30
Lathyroides	122
fativa	122
sepium	122
Vinca major	27
minor	123
Viola canina	123
hirta	123
odorata	124
odorata β .	123
tricolor; et β .	123
Viscum album	124

U.

Ulex europæus	41
Ulmus campestris	125
glabra; et β .	125
Urtica dioica	126
urens	126

E R R A T A.

The candid Reader is desired to correct these
Errors and Omissions.

Page	line
4	20 <i>after marina add β.</i>
17	9 <i>for Pluckley read Hedcorne.</i>
28	19 <i>after Sandwich add and on Shellness.</i>
36	4 <i>after Sittingbourn add and on the Standard Key.</i>
42	25 <i>after Deal add and on Shellness.</i>
49	20 <i>for elegantissimum read elegantissimum.</i>
—	22 <i>for montenum read montanum.</i>
56	22 <i>after 304 add γ.</i>
57	2 <i>after 304 add δ.</i>
59	5 <i>for uncommon read common.</i>
—	12 <i>for common read uncommon.</i>
73	4 <i>for Fairbrook read Luddenham.</i>
76	21 <i>after plentifully add on Badgen Downs, sparingly.</i>
88	14 <i>for granineum read gramineum.</i>
144	16 <i>for uncommon read common.</i>

Omitted to be inserted in their proper Places.

ANTIRRHINUM angustifolium sylvestre.

R. S. * 283.

ANTIRRHINUM Orontium. H. F. 239.

The least Snapdragon, or Calf's-snout.

In a Corn field at Luddenham — uncommon.

July. A.

CASSIDA palustris vulgatiores flore cæruleo.

R. S. 244.

SCUTELLARIA galericulata. H. F. 231.

Hooded Willow-herb.

In a moist Hedge at Chartham Hatch — uncommon.

August. P.

11-10-1964

[illegible]

Continued to be interested in their proper future.

AN FRENCH UMBRELLA

[illegible]

CASPER A. JAMES
R. S.
SOUTHERN
Hudson Valley
A new light in Southern Hudson Valley
August 19

